

The integrability of Pauli system in Lorentz violating background

To cite this article: Mariana Frank *et al*/JHEP01(2008)039

View the [article online](#) for updates and enhancements.

Related content

- [He–McKellar–Wilkins effect and scalar Aharonov–Bohm effect for a neutral particle based on the Lorentz symmetry violation](#)
K Bakke, E O Silva and H Belich
- [Hydrogen Atom and Equivalent Form of the Lévy-Leblond Equation](#)
Muhammad Adeel Ajaib
- [An analogy of the quantum hall conductivity in a Lorentz-symmetry violation setup](#)
L R Ribeiro, C Furtado and E Passos

Recent citations

- [Geometric relativistic phase from Lorentz symmetry breaking effects in the cosmic string spacetime](#)
H. Belich and K. Bakke
- [Second-Order Integrals for Systems in E²Involving Spin](#)
smet Yurduen
- [Geometric quantum phases from Lorentz symmetry breaking effects in the cosmic string spacetime](#)
H. Belich and K. Bakke

The integrability of Pauli system in Lorentz violating background

Mariana Frank and Ismail Turan

*Department of Physics, Concordia University,
7141 Sherbrooke St. West, Montreal, Quebec, H4B 1R6, Canada
E-mail: mfrank@alcor.concordia.ca, ituran@physics.concordia.ca*

İsmet Yurduşen

*Centre de Recherches Mathématiques, Université de Montréal,
CP 6128, Succ. Centre-Ville, Montréal, Québec H3C 3J7, Canada
E-mail: yurdusen@crm.umontreal.ca*

ABSTRACT: We systematically analyze the integrability of a Pauli system in Lorentz violating background at the non-relativistic level both in two- and three-dimensions. We consider the non-relativistic limit of the Dirac equation from the QED sector of the so-called Standard Model Extension by keeping only two types of background couplings, the vector a_μ and the axial vector b_μ . We show that the spin-orbit interaction comes as a higher order correction in the non-relativistic limit of the Dirac equation. Such an interaction allows the inclusion of spin degree non-trivially, and if Lorentz violating terms are allowed, they might be comparable under special circumstances. By including all possible first-order derivative terms and considering the cases $\mathbf{a} \neq 0$, $\mathbf{b} \neq 0$, and $b_0 \neq 0$ one at a time, we determine the possible forms of constants of motion operator, and discuss the existence or continuity of integrability due to Lorentz violating background.

KEYWORDS: Discrete and Finite Symmetries, Integrable Equations in Physics.

Contents

1. Introduction	1
2. The model and its non-relativistic limit	3
3. The analysis of 2D Pauli system in LV background	5
3.1 Lorentz violation with vector coupling ($a_i \neq 0$):	5
3.2 Lorentz violation with pure spacelike axial-vector coupling ($b_i \neq 0$):	7
3.3 Lorentz violation with pure timelike axial-vector coupling ($b_0 \neq 0$):	8
4. The analysis of 3D Pauli system in LV background	9
4.1 Lorentz violation with vector coupling ($a_i \neq 0$):	12
4.2 Lorentz violation with pure spacelike axial-vector coupling ($b_i \neq 0$):	13
4.3 Lorentz violation with pure timelike axial-vector coupling ($b_0 \neq 0$):	13
5. Perturbative solutions	14
6. Conclusion	15

1. Introduction

In classical mechanics, integrability of a Hamiltonian system can be defined as the existence of a set of n functionally independent constants of motion X_1, X_2, \dots, X_n including the Hamiltonian itself if the system has n degrees of freedom. These constants of motion must be in involution. The concept of integrability can be extended to so-called superintegrability which requires at least one additional constant of motion (minimally superintegrable) and allows totally $n-1$ additional constants of motion (maximally superintegrable).¹ These additional constants of motion, however, are not necessarily in involution among each other, nor with X_1, X_2, \dots, X_{n-1} . In a very similar manner, the notion of (super)integrability can also be defined in quantum mechanics through well-defined linear constants of motion operators which are now supposed to be algebraically independent [1–4]. In quantum mechanics, integrability not only simplifies the calculation of energy levels and wave functions, but also provides a complete set of quantum numbers, which characterize the system completely. Superintegrability, on the other hand, may entail exact solvability. The harmonic oscillator [2], and the Kepler, or Coulomb system [1] are well known examples of superintegrability.

¹This is due to the fact that the phase space is $2n$ dimensional and at least one free degree of freedom has to be left in order to have dynamics in the system.

Systematic searching for superintegrability in Hamiltonian systems with kinetic energy quadratic in momenta and with only coordinate dependent potentials was started quite some time ago [3]. In these works the constants of motion were considered up to second-order in momenta in order to search for quadratic (super)integrability. Both in classical and quantum mechanics, second-order integrability of such Hamiltonians mimics the separations of variables which is the first step for exact solvability. Such close connection doesn't hold for the systems with velocity dependent potentials $V(\mathbf{p}, \mathbf{A})$ [5].

Recently, the (super)integrability of spin-dependent Hamiltonian systems with a generic scalar potential has been studied in a systematic way up to first-order [6]. There are other known systems which introduce velocity-dependent interactions without requiring an external vector potential \mathbf{A} . For example, if one allows violation of Lorentz symmetry at the relativistic level, the extra fields which describe such violation can also induce velocity-dependent interactions at non-relativistic level. The aim of this study is to systematically analyze (super)integrability of a Hamiltonian system (more precisely the hydrogen atom) with spin-orbit interactions in some Lorentz violating backgrounds both in 2- and 3-dimensions.

There exists a well established framework [7], called Standard-Model Extension (SME), for the study of the Lorentz and Charge-Parity-Time reversal (CPT) violation. The Standard Model Extension is the generalization of the Standard Model (SM) with additional Lorentz and CPT violating interactions introduced through some tensorial background fields. It can be considered a low-energy limit of a fundamental theory in which the Lorentz and CPT symmetries are exact but broken spontaneously when evolved down to low energy scale, due to the existence of these tensorial background fields. SME is one of the elegant way to formulate the problem. Of course the coordinate independence of physical observables are maintained by requiring *observer* Lorentz invariance which describes the transformations of coordinates. Lorentz violation considered in the model is so called *particle* Lorentz violation which describes rotations and boosts of particles and localized fields but not background fields in a specific observer's inertial frame. Formulating the model in non-Minkowski spacetimes [8] leads to spacetime dependent coupling coefficients. Our approach here is minimalistic, i.e., constant coefficients.

In this study we consider only the effects of vectorial background couplings a_μ and b_μ to the (super)integrability of what we call the Pauli system. Here a_μ and b_μ are related to the vacuum expectation values of some vectorial background fields. Seeking (super)integrability for a Pauli system in such LV background and aiming exact solvability could be of some interest under certain circumstances. For example, the current bound on b_0 [9], the time component of b_μ , is $b_0^{-1} \gtrsim 10^{-3} \text{ cm}$ and it is much weaker than that on $|\mathbf{b}|$, the space part of the b_μ . So, for a system with timelike b_μ coupling the perturbative approach fails if the effective size of the system is of the order of 10^{-3} cm , and then the search for an exact solution becomes unavoidable.

The outline of the paper is as follows. In the next section, section 2, we introduce the relevant part of the SME Lagrangian and consider the non-relativistic limit. Then, in section 3, the 2-dimensional Pauli system in LV background is discussed in three separate cases. In section 4, before discussing the generalization of the problem to 3-dimensions,

we present the determining equations in the most general form. We then shortly discuss the perturbative approaches to the system in section 5 and conclude in section 6.

2. The model and its non-relativistic limit

There exist vast number of studies on the SME both at theoretical and phenomenological levels [10, 11]. Here we work in the QED sector of the SME for electron [12–14]. The QED Lagrangian for electron can be given as

$$\begin{aligned} \mathcal{L}_{\text{electron}} &= \frac{i}{2} \bar{\psi} \Gamma^\mu \overleftrightarrow{D}_\mu \psi - \bar{\psi} M \psi, \\ \Gamma_\mu &= \gamma_\mu + c_{\lambda\mu} \gamma^\lambda + d_{\lambda\mu} \gamma_5 \gamma^\lambda + \dots \\ M &= m + a_\mu \gamma^\mu + b_\mu \gamma_5 \gamma^\mu + \frac{1}{2} H_{\mu\nu} \sigma^{\mu\nu}, \end{aligned} \tag{2.1}$$

where $D_\mu = \partial_\mu + ieA_\mu$ with the vector potential A_μ ($f \overleftrightarrow{D}_\mu g \equiv f D_\mu g - (D_\mu f) g$), a_μ and b_μ are the (pseudo-)vectorial CPT -odd coefficients and the other terms are CPT -even tensorial coefficients. These are the only terms that are obtained from the SME but Γ_μ can have more terms originating from non-renormalizable higher-dimensional operators.

It is clear from the above Lagrangian that LV doesn't require CPT violation but the vice versa is true [15]. In this study, we consider the only two CPT -odd vectorial couplings a_μ and b_μ and neglect the tensorial ones.² A comprehensive analysis of field redefinitions and redundant parameters of the model can be found in ref. [16]. For example, the a_μ term can be absorbed by redefining the fermion field by a phase factor $\exp(ia_\mu x^\mu)$ and thus the spectrum will be unaffected by such term. Depending on the complexity of the model (photon interaction, fermion mixings, etc.), this situation could change. A similar transformation can be found for the b_μ case if, for example, a free massless fermion is assumed, but in general the b_μ term is non-trivial. So, in our discussion we keep both a_μ and b_μ terms.

We now consider the non-relativistic limit of the Dirac equation for electron starting from eq. (2.1) with LV vectorial couplings only. We take the external electromagnetic field as $A_\mu = (A_0, \mathbf{0})$. The Dirac equation becomes³

$$(p_\mu \gamma^\mu - eA_\mu \gamma^\mu - a_\mu \gamma^\mu - \gamma_5 b_\mu \gamma^\mu - m) \psi = 0, \tag{2.2}$$

where we adopt the Dirac representation for the γ matrices. For convenience one can define the energy of the electron $p_0 = p'_0 + m$ (by extracting the rest energy) so that p'_0 becomes much smaller than m in the non-relativistic limit. If we also assume $eA_0, |a|, |b| \ll m$, we obtain the following equation for the 2×1 spinor ψ_1 , which is the large component of ψ ,

$$\begin{aligned} \left[(p'_0 - m - a_0 - \boldsymbol{\sigma} \cdot \mathbf{b}) - \frac{1}{2m} (\boldsymbol{\sigma} \cdot (\mathbf{p} - \mathbf{a}) - b_0) \Lambda^{-1} (\boldsymbol{\sigma} \cdot (\mathbf{p} - \mathbf{a}) - b_0) \right] \psi_1 = 0, \\ \left(1 + \frac{p'_0 - eA_0 - a_0 - \boldsymbol{\sigma} \cdot \mathbf{b}}{2m} \right) \equiv \Lambda, \end{aligned} \tag{2.3}$$

²Note that allowing non-renormalizable higher-dimensional operators induces additional vectorial fields.

³We use the natural units, $\hbar = 1, c = 1$, throughout the paper.

	Rotation	C	P	T	CPT
$\mathbf{a} \cdot \mathbf{p}$	-	-	-	-	-
$\boldsymbol{\sigma} \cdot \mathbf{b}$	-	+	+	-	-
$b_0 \boldsymbol{\sigma} \cdot \mathbf{p}$	+	+	-	+	-

Table 1: Properties of LV terms in the Hamiltonian eq. (2.5) under C , P , T , and rotational symmetries.

where $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ are the Pauli matrices. Keeping the first two terms in Λ^{-1} and making some straightforward manipulations lead to the following equation

$$\begin{aligned}
 i\frac{\partial}{\partial t}\psi_1 &= H_{\text{NR}}\psi_1, \\
 H_{\text{NR}} &= \frac{\mathbf{p}^2}{2m} + eA_0 - \frac{e}{4m^2}\boldsymbol{\sigma} \cdot \mathbf{E} \times \mathbf{p} - \frac{\mathbf{p}^4}{8m^3} + \frac{ie}{4m^2}\mathbf{E} \cdot \mathbf{p} + H_{\text{LV}}, \\
 H_{\text{LV}} &= -\frac{1}{m}\mathbf{a} \cdot \mathbf{p} + \boldsymbol{\sigma} \cdot \mathbf{b} - \frac{b_0}{m}\boldsymbol{\sigma} \cdot \mathbf{p} + \frac{a_0}{4m^2}\mathbf{p}^2 + \frac{1}{4m^2}\boldsymbol{\sigma} \cdot \mathbf{p} \boldsymbol{\sigma} \cdot \mathbf{b}, \quad (2.4)
 \end{aligned}$$

where \mathbf{E} is the electric field and we keep up to linear terms in LV parameters, since they are presumed small, and we also neglect the constant terms. In H_{NR} , the first order correction terms to the usual Hamiltonian are the spin-orbit, relativistic, and the potential energy correction, respectively. We assume that the potential energy is spherically symmetric, $V_0(r) \equiv eA_0(r)$ so that the term $\boldsymbol{\sigma} \cdot \mathbf{E} \times \mathbf{p}$ with $\mathbf{E} = \frac{1}{r}\frac{dA_0}{dr}\mathbf{r}$ leads to the usual spin-orbit interaction $\boldsymbol{\sigma} \cdot \mathbf{L}$ with \mathbf{L} the angular momentum operator.

Had we neglected all these three corrections and considered the Lorentz symmetry preserved, there would be no spin dependence in the system (since we set the external magnetic field zero) and the spin degree of the electron would become trivial. The only spin dependence is through $\boldsymbol{\sigma} \cdot \mathbf{L}$ and we intend to keep that term and neglect the other two whose presence would considerably complicate our study. As seen from eq. (2.4), H_{LV} also brings spin dependent interactions only through b_μ . The last two terms of H_{LV} come from the second term in Λ^{-1} expansion and are higher order with respect to the first three terms of H_{LV} . Hence, we neglect them in the rest of our calculation. We further set $m = 1$ for simplicity.

Taking into account all of the above remarks, and inserting the Coulomb potential of the nucleus, the Lorentz violating Hamiltonian H_{NR} for the hydrogen atom follows from eq. (2.4) as

$$H = \frac{1}{2}(p_1^2 + p_2^2 + p_3^2) + V_0(r) + V_1(r)\boldsymbol{\sigma} \cdot \mathbf{L} - \mathbf{a} \cdot \mathbf{p} + \boldsymbol{\sigma} \cdot \mathbf{b} - b_0 \boldsymbol{\sigma} \cdot \mathbf{p}, \quad (2.5)$$

where $V_0 = \Gamma_1/r$, $V_1 = \Gamma_2/r^3$ with $\Gamma_1 = -e^2$ and $\Gamma_2 = -e^2/4$. In the absence of the LV terms we call the above Hamiltonian the Pauli system, even though a system with an external magnetic field is usually considered as Pauli system. The properties of the LV terms under the Charge (C), Parity (P), Time-reversal (T), and rotational symmetries are given in table 1.

In order to investigate the first-order integrability and/or superintegrability we analyze the commutativity of a general first-order integral of motion⁴

$$X = (A_0 + \mathbf{A} \cdot \boldsymbol{\sigma})p_1 + (B_0 + \mathbf{B} \cdot \boldsymbol{\sigma})p_2 + (C_0 + \mathbf{C} \cdot \boldsymbol{\sigma})p_3 + \Phi_0 + \boldsymbol{\Phi} \cdot \boldsymbol{\sigma} - \frac{i}{2} \{ (A_0 + \mathbf{A} \cdot \boldsymbol{\sigma})_x + (B_0 + \mathbf{B} \cdot \boldsymbol{\sigma})_y + (C_0 + \mathbf{C} \cdot \boldsymbol{\sigma})_z \}, \quad (2.6)$$

with the Hamiltonian given in eq. (2.5). In eq. (2.6), \mathbf{A} , \mathbf{B} , \mathbf{C} and $\boldsymbol{\Phi}$ where $\mathbf{A} = (A_1, A_2, A_3)$ (\mathbf{B} , \mathbf{C} and $\boldsymbol{\Phi}$ are defined similarly) are real functions of \mathbf{r} . The commutativity relation

$$[H, X] = 0, \quad (2.7)$$

has second-, first-, and zeroth-order terms in the momenta. Setting the coefficients of each of these terms equal to zero we get equations determining A_0 , B_0 , C_0 , Φ_0 and A_i , B_i , C_i , Φ_i ($i = 1, 2, 3$).

3. The analysis of 2D Pauli system in LV background

In this section we consider the integrability of 2D Pauli system in LV background. There are three LV terms in the Hamiltonian given in eq. (2.5). For the sake of simplicity we consider the effects of each term one at a time; the case with non-zero vector coupling ($a_i \neq 0$), with pure spacelike axial-vector coupling ($b_i \neq 0$), and finally with pure timelike axial-vector coupling ($b_0 \neq 0$).

3.1 Lorentz violation with vector coupling ($a_i \neq 0$):

In the absence of the axial-vector coupling terms in the Hamiltonian, eq. (2.5) in two-dimensions is given as

$$H_{2D}^{(a_\mu \neq 0)} = \frac{1}{2} (p_1^2 + p_2^2) + V_0(\rho) + V_1(\rho) \sigma_3 L_3 - a_1 p_1 - a_2 p_2, \quad (3.1)$$

where again $V_0 = \Gamma_1/\rho$ and $V_1 = \Gamma_2/\rho^3$, Γ_1 and Γ_2 are constants defined in the previous section. Since now the Hamiltonian eq. (3.1) is a diagonal matrix we can also choose X given in eq. (2.6) as diagonal too and write

$$X_{2D}^{(a_\mu \neq 0)} = (A_0 + A_3 \sigma_3)p_1 + (B_0 + B_3 \sigma_3)p_2 + \Phi_0 + \Phi_3 \sigma_3 - \frac{i}{2} \{ (A_0 + A_3 \sigma_3)_x + (B_0 + B_3 \sigma_3)_y \}. \quad (3.2)$$

The requirement of vanishing of the commutator $[H_{2D}^{(a_\mu \neq 0)}, X_{2D}^{(a_\mu \neq 0)}]$ gives us a total of 12 equations for A_ν , B_ν and Φ_ν , ($\nu = 0, 3$). The technique is to start from the highest order terms (second-order in our problem) and determine some functions in $X_{2D}^{(a_\mu \neq 0)}$ and then use these solutions to apply the same procedure repeatedly for the remaining orders until

⁴Throughout the paper, the subscripts x , y and z represent the partial derivatives with respect to the Cartesian coordinates.

all unknown functions in $X_{2D}^{(a_\mu \neq 0)}$ are fixed. So, let's start from the second-order. We get the following six partial differential equations

$$A_{\nu x} = 0, \quad B_{\nu y} = 0, \quad A_{\nu y} + B_{\nu x} = 0, \quad (\nu = 0, 3), \quad (3.3)$$

which can immediately be integrated to give

$$A_\nu = \xi_\nu + w_\nu y, \quad B_\nu = \eta_\nu - w_\nu x. \quad (3.4)$$

After introducing eq. (3.4) into the coefficients of the first-order terms and separating the real and imaginary parts we have 4 determining equations for Φ_ν

$$\begin{aligned} \Phi_{\nu x} &= \delta_{\nu,3-\kappa}[(w_\kappa x - \eta_\kappa)yV_{1y} - (w_\kappa y + \xi_\kappa)yV_{1x} - \eta_\kappa V_1] + a_2 w_\nu, \\ \Phi_{\nu y} &= \delta_{\nu,3-\kappa}[(w_\kappa y + \xi_\kappa)xV_{1x} - (w_\kappa x - \eta_\kappa)xV_{1y} + \xi_\kappa V_1] - a_1 w_\nu, \quad (\nu = 0, 3), \end{aligned} \quad (3.5)$$

where $\delta_{\nu,\kappa}$ is the Kronecker delta function. In order to satisfy the compatibility conditions for Φ_ν we must have $\xi_\nu = 0$ and $\eta_\nu = 0$, $\nu = 0, 3$. Then, eq. (3.5) can be easily integrated to give

$$\Phi_\nu = w_\nu(a_2 x - a_1 y), \quad (\nu = 0, 3), \quad (3.6)$$

where we used $V_1 = \Gamma_2/\rho^3$ and set the integration constants equal to zero without loss of generality.

Two of the 12 determining equations, which are indeed the coefficients of the zeroth-order terms, have not been used up to now. If we introduce all the information gathered from the coefficients of the higher-order terms into the coefficients of the zeroth-order terms we get the following two conditions to be satisfied

$$(a_1 x + a_2 y)V_1 w_\nu = 0, \quad (\nu = 0, 3). \quad (3.7)$$

Clearly we must have either (1) both $a_1 = 0$ and $a_2 = 0$, the components of the LV vector a_μ on the plane or (2) $w_\nu = 0$.

Case (1) corresponds to a LV vector of the form $a_\mu = (a_0, 0, 0, a_3)$ or even a pure timelike vector ($a_3 = 0$). The latter is trivial since the system still remains Pauli system up to some constants. The former is non-trivial since a non-zero space component a_3 is allowed but it doesn't couple with the electron in xy plane. Therefore, under these conditions only, the symmetries of the Pauli system are restored. For example, we get the constant of motion as

$$\begin{aligned} X_{2D}^{(a_\mu \neq 0)}(a_1 = 0 = a_2) &= -(w_0 + w_3 \sigma_3)(xp_2 - yp_1) \\ &= -(w_0 + w_3 \sigma_3)L_3, \end{aligned} \quad (3.8)$$

where w_0 and w_3 are arbitrary constants. As expected both $L_3 \otimes I$ and $L_3 \otimes \sigma_3$ are constants of motion. However, they are not different from each other since σ_3 trivially commutes with the Hamiltonian in 2D. Therefore, there is only one extra constant of motion other than the Hamiltonian and there exists first-order integrability.

For the Case (2), which is indeed the non-trivial one, $X_{2D}^{(a_\mu \neq 0)}$ vanishes as seen from eq. (3.8) and there is no first-order integrability.

3.2 Lorentz violation with pure spacelike axial-vector coupling ($b_i \neq 0$):

If we only have pure spacelike axial-vector coupling, i.e., $b_i \neq 0$, then in two-dimensions the Hamiltonian eq. (2.5) and the integral of motion eq. (2.6) are given as

$$H_{2D}^{(b_i \neq 0)} = \frac{1}{2}(p_1^2 + p_2^2) + V_0(\rho) + V_1(\rho) \sigma_3 L_3 + \boldsymbol{\sigma} \cdot \mathbf{b}, \quad (3.9)$$

$$X_{2D}^{(b_i \neq 0)} = (A_0 + \mathbf{A} \cdot \boldsymbol{\sigma})p_1 + (B_0 + \mathbf{B} \cdot \boldsymbol{\sigma})p_2 + \Phi_0 + \boldsymbol{\Phi} \cdot \boldsymbol{\sigma} - \frac{i}{2} \{ (A_0 + \mathbf{A} \cdot \boldsymbol{\sigma})_x + (B_0 + \mathbf{B} \cdot \boldsymbol{\sigma})_y \}, \quad (3.10)$$

where $V_0 = \Gamma_1/\rho$ and $V_1 = \Gamma_2/\rho^3$ are assumed. Since the Hamiltonian eq. (3.9) is not diagonal, we can no longer choose the integral of motion operator $X_{2D}^{(b_i \neq 0)}$ diagonal.

In this case, unlike the previous case, there are 12 determining equations coming from the coefficients of the second-order terms since we have non-diagonal entries which double the number of equations. The ones from the diagonal elements are exactly the same as eq. (3.3) and hence the solutions for A_ν and B_ν , ($\nu = 0, 3$) as given in eq. (3.4), are the same. From the off-diagonal elements we also have the following 6 equations

$$\begin{aligned} 2yA_1V_1 + A_{2x} &= 0, & 2yA_2V_1 - A_{1x} &= 0, \\ 2xB_2V_1 + B_{1y} &= 0, & 2xB_1V_1 - B_{2y} &= 0, \\ 2V_1(xA_2 - yB_2) + A_{1y} + B_{1x} &= 0, & 2V_1(xA_1 - yB_1) - A_{2y} + B_{2x} &= 0. \end{aligned} \quad (3.11)$$

However, there is only a so called *trivial solution* (keeping in mind that $V_1 = \Gamma_2/\rho^3$) for the system of equations given in eq. (3.11), which is $A_1 = A_2 = B_1 = B_2 = 0$. This has been shown both by hand and by using the Maple software [17].

When we introduce all of this information into the rest of the determining equations we get, in addition to the equations given in eq. (3.5) (with of course $a_1 = 0$ and $a_2 = 0$), the following four equations for the contribution of the coefficients of the first-order terms

$$\begin{aligned} 2b_2w_3y + 2yV_1\Phi_2 - \Phi_{1x} &= 0, & 2b_1w_3y + 2yV_1\Phi_1 + \Phi_{2x} &= 0, \\ 2b_2w_3x + 2xV_1\Phi_2 + \Phi_{1y} &= 0, & 2b_1w_3x + 2xV_1\Phi_1 - \Phi_{2y} &= 0. \end{aligned} \quad (3.12)$$

These equations can be solved and we find that $\xi_\nu = 0$, $\eta_\nu = 0$, $\Phi_0 = 0$ ($\nu = 0, 3$) are solutions if either of the following conditions is satisfied

- Case (1): $b_1 = 0 = b_2$
- Case (2): $w_3 = 0$.

Note that the determining equations coming from the coefficients of the zeroth-order terms are identically satisfied in either case. The constant of motion operator is the same as given in eq. (3.6) but, since this time there is no condition on w_0 , $L_3 \otimes I$ becomes a constant of motion. It is also important to note here that since the Hamiltonian eq. (3.9) has σ_1 and σ_2 in it, the fact that σ_3 commutes with the Hamiltonian is no longer *trivial*. Hence, the term $L_3 \otimes \sigma_3$ that comes with w_3 should now be considered as independent. Obviously, in

the presence of b_1 and b_2 (Case (2)) we must have $w_3 = 0$ which leaves $L_3 \otimes I$ as the only constant of motion. However, Case (1) puts no constraint on the z-component of b_μ , and if the coupling vector is perpendicular to the motion of the plane ($b_z \neq 0$), both $L_3 \otimes I$ and $L_3 \otimes \sigma_3$ are constants of motion but of course they are not independent. Therefore, including the Hamiltonian there are two constants of motion, independent of which of the above cases are satisfied. The first-order integrability is preserved in the presence of a LV background \mathbf{b} with even arbitrary nonzero components in each direction.

3.3 Lorentz violation with pure timelike axial-vector coupling ($b_0 \neq 0$):

When $\mathbf{a} = \mathbf{0}$, $\mathbf{b} = \mathbf{0}$ but $b_0 \neq 0$ the Hamiltonian eq. (2.5) and the integral of motion eq. (2.6) reduce to the following in two-dimensions

$$H_{2D}^{(b_0 \neq 0)} = \frac{1}{2}(p_1^2 + p_2^2) + V_0(\rho) + V_1(\rho) \sigma_3 L_3 - b_0(\sigma_1 p_1 + \sigma_2 p_2), \quad (3.13)$$

$$X_{2D}^{(b_0 \neq 0)} = (A_0 + \mathbf{A} \cdot \boldsymbol{\sigma}) p_1 + (B_0 + \mathbf{B} \cdot \boldsymbol{\sigma}) p_2 + \Phi_0 + \boldsymbol{\Phi} \cdot \boldsymbol{\sigma} - \frac{i}{2} \{ (A_0 + \mathbf{A} \cdot \boldsymbol{\sigma})_x + (B_0 + \mathbf{B} \cdot \boldsymbol{\sigma})_y \}, \quad (3.14)$$

where $V_0 = \Gamma_1/\rho$ and $V_1 = \Gamma_2/\rho^3$ are assumed. Again, because of the fact that the Hamiltonian eq. (3.13) is not diagonal, we can no longer choose the integral of motion diagonal as in the previous case.

In this case we have 12 equations coming from the coefficients of the second-order terms. Three of them are exactly the same with eq. (3.3) for $\nu = 0$ and hence we have the same A_0 and B_0 as in eq. (3.4). The rest of the determining equations from the coefficients of the second-order terms can be written as

$$\begin{aligned} 2b_0 A_2 + A_{3x} = 0, \quad 2b_0 B_1 - B_{3y} = 0, \quad 2y A_2 V_1 - A_{1x} = 0, \quad 2x B_1 V_1 - B_{2y} = 0, \\ 2b_0 A_3 - 2y A_1 V_1 - A_{2x} = 0, \quad 2b_0 B_3 + 2x B_2 V_1 + B_{1y} = 0, \quad 2b_0 A_1 - 2b_0 B_2 - A_{3y} - B_{3x} = 0, \\ 2b_0 A_3 + 2V_1(x A_2 - y B_2) + A_{1y} + B_{1x} = 0, \quad 2b_0 B_3 + 2V_1(x A_1 - y B_1) - A_{2y} - B_{2x} = 0. \end{aligned} \quad (3.15)$$

There is only a so called trivial solution for the system of equations given in eq. (3.15), which is $\mathbf{A} = \mathbf{0}$ and $\mathbf{B} = \mathbf{0}$. This is shown again by using the Maple software [17].

After introducing the already found functions A_0 , B_0 , \mathbf{A} , and \mathbf{B} into the determining equations coming from the coefficients of the first-order terms we have the following 8 partial differential equations for Φ_0 and $\boldsymbol{\Phi}$

$$\begin{aligned} \Phi_{0x} = 0, \quad \Phi_{0y} = 0, \\ \Phi_{1x} = 2y V_1 \Phi_2, \quad \Phi_{1y} = -2x V_1 \Phi_2 - b_0(w_0 + 2\Phi_3), \\ \Phi_{2x} = -2y V_1 \Phi_1 + b_0(w_0 + 2\Phi_3), \quad \Phi_{2y} = 2x V_1 \Phi_1, \\ \Phi_{3x} = -y(w_0 y + \xi_0) V_{1x} + y(w_0 x - \eta_0) V_{1y} - \eta_0 V_1 - 2b_0 \Phi_2, \\ \Phi_{3y} = x(w_0 y + \xi_0) V_{1x} - x(w_0 x - \eta_0) V_{1y} + \xi_0 V_1 + 2b_0 \Phi_1. \end{aligned} \quad (3.16)$$

It is immediately seen that $\Phi_0 = \text{const}$, which can be taken as zero without loss of generality. Then we are left with 6 first-order partial differential for $\boldsymbol{\Phi}$. The requirement of the equality

of the mixed partial derivatives give 3 more equations for Φ and their first-order derivatives. Now, if we eliminate the first-order derivatives of Φ in this system by using the equations given in eq. (3.16) we get a system of algebraic equations for Φ . In order to solve this system the rank of the coefficient matrix has to be equal to the rank of the extended matrix, determined by adding the non homogeneous vector as a column. A simple analysis shows that this condition can only be possible if $\xi_0 = 0$ and $\eta_0 = 0$. In this case we have the following solution

$$\Phi_1 = -\frac{y}{x}\Phi_2, \quad \Phi_3 = -\frac{w_0}{2} + \left(\frac{1}{2b_0x} + \frac{b_0}{V_{1,x}}\right)\Phi_2. \quad (3.17)$$

Finally, we get the following 4 partial differential equations from the coefficients of the zeroth-order terms

$$\begin{aligned} 2b_0\Phi_{3x} - 2V_1(y\Phi_{1x} - x\Phi_{1y}) - (\Phi_{2xx} + \Phi_{2yy}) &= 0, & 2b_0(\Phi_{1y} - \Phi_{2x}) - (\Phi_{3xx} + \Phi_{3yy}) &= 0, \\ 2b_0\Phi_{3y} - 2V_1(y\Phi_{2x} - x\Phi_{2y}) + (\Phi_{1xx} + \Phi_{1yy}) &= 0, & y\Phi_2 + x(x\Phi_{2y} - y\Phi_{2x}) &= 0, \end{aligned} \quad (3.18)$$

where $\mathbf{A} = \mathbf{0}$, $\mathbf{B} = \mathbf{0}$, $A_0 = w_0 y$ and $B_0 = -w_0 x$ are used. Together with eq. (3.17) it is possible to show that only $\Phi_2 = 0$ satisfies simultaneously all the 4 differential equations in eq. (3.18). Thus, to sum up we find the following set of solutions in the case if $b_0 \neq 0$:

$$\begin{aligned} A_0 = w_0 y, & & B_0 = -w_0 x, & & \mathbf{A} = \mathbf{0}, & & \mathbf{B} = \mathbf{0}, \\ \Phi_0 = 0, & & \Phi_1 = 0, & & \Phi_2 = 0, & & \Phi_3 = -w_0/2. \end{aligned} \quad (3.19)$$

It is seen from the constant of motion operator in eq. (3.14) that we have

$$\begin{aligned} X_{2D}^{(b_0 \neq 0)} &= w_0(y p_1 - x p_2) - \sigma_3 w_0/2 \\ &= -w_0(L_3 + \sigma_3/2) \end{aligned} \quad (3.20)$$

which is nothing but the total angular momentum J_3 . It is important to note here that, as opposed to the case in 2D where there are no LV terms and hence both L_3 and J_3 are constants of motion, only J_3 is a constant of motion in this case ($b_0 \neq 0$). The fact that L_3 does not commute with the Hamiltonian is something expected from the form of the LV term in eq. (3.13). However, first-order integrability is still restored.

4. The analysis of 3D Pauli system in LV background

In general in the presence of LV terms we have the Hamiltonian and integral of motion given in equations eq. (2.5) and eq. (2.6) respectively in 3-dimensional Euclidean spaces. In a similar fashion as in the two-dimensional case it is also convenient to analyze the problem in three separate cases. However, it is advantageous to give the determining equations coming from the coefficients of the second- and first-order terms in full generality (that is, keeping \mathbf{a} , \mathbf{b} , and b_0 non zero) before analyzing them in case by case.

i) Determining equations coming from the second-order terms: From the diagonal elements it is immediately found that we have the following 6 determining equations

$$\begin{aligned} A_{0x} &= 0, & B_{0y} &= 0, & C_{0z} &= 0, \\ A_{0y} + B_{0x} &= 0, & A_{0z} + C_{0x} &= 0, & B_{0z} + C_{0y} &= 0, \end{aligned} \quad (4.1)$$

from which the following solutions are obvious

$$\begin{aligned} A_0 &= \beta_1 - \alpha_3 y + \alpha_2 z, \\ B_0 &= \beta_2 + \alpha_3 x - \alpha_1 z, \\ C_0 &= \beta_3 - \alpha_2 x + \alpha_1 y. \end{aligned} \quad (4.2)$$

where α_i and β_i ($i = 1, 2, 3$) are integration constants. Note that these solutions are independent of the LV background couplings so that they hold for each of the cases discussed below. If we compare eq. (4.2) with eq. (3.4) we see that we have the following correspondences: $\beta_1 = \xi_0$, $\beta_2 = \eta_0$ and $\alpha_3 = -w_0$. After introducing the equation (4.2) into the rest of the coefficients of the second-order terms and separating the imaginary and real parts of the coefficients coming from the off-diagonal elements we are left with an over determined system of 18 partial differential equations for A_i, B_i, C_i ($i = 1, 2, 3$). These are,

$$2zA_1V_1 + A_{3x} + 2b_0A_2 = 0, \quad (4.3)$$

$$2yA_1V_1 + A_{2x} - 2b_0A_3 = 0, \quad (4.4)$$

$$2xB_2V_1 + B_{1y} + 2b_0B_3 = 0, \quad (4.5)$$

$$2zB_2V_1 + B_{3y} - 2b_0B_1 = 0, \quad (4.6)$$

$$2xC_3V_1 + C_{1z} - 2b_0C_2 = 0, \quad (4.7)$$

$$2yC_3V_1 + C_{2z} + 2b_0C_1 = 0, \quad (4.8)$$

$$2V_1(yA_2 + zA_3) - A_{1x} = 0, \quad (4.9)$$

$$2V_1(xB_1 + zB_3) - B_{2y} = 0, \quad (4.10)$$

$$2V_1(xC_1 + yC_2) - C_{3z} = 0, \quad (4.11)$$

$$2zV_1(A_2 + B_1) + A_{3y} + B_{3x} + 2b_0(B_2 - A_1) = 0, \quad (4.12)$$

$$2yV_1(A_3 + C_1) + A_{2z} + C_{2x} + 2b_0(A_1 - C_3) = 0, \quad (4.13)$$

$$2xV_1(B_3 + C_2) + B_{1z} + C_{1y} + 2b_0(C_3 - B_2) = 0, \quad (4.14)$$

$$2V_1(xA_1 + yA_2 - zC_1) - A_{3z} - C_{3x} - 2b_0C_2 = 0, \quad (4.15)$$

$$2V_1(xB_1 + yB_2 - zC_2) - B_{3z} - C_{3y} + 2b_0C_1 = 0, \quad (4.16)$$

$$2V_1(xA_2 - yB_2 - zB_3) + A_{1y} + B_{1x} + 2b_0A_3 = 0, \quad (4.17)$$

$$2V_1(xA_1 + zA_3 - yB_1) - A_{2y} - B_{2x} + 2b_0B_3 = 0, \quad (4.18)$$

$$2V_1(xA_3 - yC_2 - zC_3) + A_{1z} + C_{1x} - 2b_0A_2 = 0, \quad (4.19)$$

$$2V_1(yB_3 - xC_1 - zC_3) + B_{2z} + C_{2y} + 2b_0B_1 = 0. \quad (4.20)$$

ii) Determining equations coming from the first-order terms: We set the coefficients of the p_i ($i = 1, 2, 3$) zero at each entry of the commutation relation. After introducing equations (4.2) and separating the real and imaginary parts, we have the following

21 partial differential equations, 12 of which are

$$x \hat{\mathbf{r}} \cdot \boldsymbol{\beta} \dot{V}_1 + (\beta_1 - y\alpha_3 + 2y\Phi_3)V_1 + \Phi_{2z} - \mathbf{a} \cdot \nabla C_2 - b_0(\alpha_1 - 2\Phi_1) - 2(\mathbf{b} \times \mathbf{C})_2 = 0, \quad (4.21)$$

$$x \hat{\mathbf{r}} \cdot \boldsymbol{\beta} \dot{V}_1 + (\beta_1 + z\alpha_2 - 2z\Phi_2)V_1 - \Phi_{3y} + \mathbf{a} \cdot \nabla B_3 - b_0(\alpha_1 - 2\Phi_1) + 2(\mathbf{b} \times \mathbf{B})_3 = 0, \quad (4.22)$$

$$y \hat{\mathbf{r}} \cdot \boldsymbol{\beta} \dot{V}_1 + (\beta_2 - z\alpha_1 + 2z\Phi_1)V_1 + \Phi_{3x} - \mathbf{a} \cdot \nabla A_3 - b_0(\alpha_2 - 2\Phi_2) - 2(\mathbf{b} \times \mathbf{A})_3 = 0, \quad (4.23)$$

$$y \hat{\mathbf{r}} \cdot \boldsymbol{\beta} \dot{V}_1 + (\beta_2 + x\alpha_3 - 2x\Phi_3)V_1 - \Phi_{1z} + \mathbf{a} \cdot \nabla C_1 - b_0(\alpha_2 - 2\Phi_2) + 2(\mathbf{b} \times \mathbf{C})_1 = 0, \quad (4.24)$$

$$z \hat{\mathbf{r}} \cdot \boldsymbol{\beta} \dot{V}_1 + (\beta_3 - x\alpha_2 + 2x\Phi_2)V_1 + \Phi_{1y} - \mathbf{a} \cdot \nabla B_1 - b_0(\alpha_3 - 2\Phi_3) - 2(\mathbf{b} \times \mathbf{B})_1 = 0, \quad (4.25)$$

$$z \hat{\mathbf{r}} \cdot \boldsymbol{\beta} \dot{V}_1 + (\beta_3 + y\alpha_1 - 2y\Phi_1)V_1 - \Phi_{2x} + \mathbf{a} \cdot \nabla A_2 - b_0(\alpha_3 - 2\Phi_3) + 2(\mathbf{b} \times \mathbf{A})_2 = 0, \quad (4.26)$$

$$(y\alpha_2 + z\alpha_3 - 2y\Phi_2 - 2z\Phi_3)V_1 + \Phi_{1x} - \mathbf{a} \cdot \nabla A_1 - 2(\mathbf{b} \times \mathbf{A})_1 = 0, \quad (4.27)$$

$$(x\alpha_1 + z\alpha_3 - 2x\Phi_1 - 2z\Phi_3)V_1 + \Phi_{2y} - \mathbf{a} \cdot \nabla B_2 - 2(\mathbf{b} \times \mathbf{B})_2 = 0, \quad (4.28)$$

$$(x\alpha_1 + y\alpha_2 - 2x\Phi_1 - 2y\Phi_2)V_1 + \Phi_{3z} - \mathbf{a} \cdot \nabla C_3 - 2(\mathbf{b} \times \mathbf{C})_3 = 0, \quad (4.29)$$

$$\begin{aligned} \Phi_{0x} = & \left((yA_{3x} - xA_{3y}) + (xA_{2z} - zA_{2x}) + (zA_{1y} - yA_{1z}) + (C_2 - B_3) \right) V_1 \\ & - \left(x(\hat{\mathbf{r}} \times \mathbf{A})_1 + y(\hat{\mathbf{r}} \times \mathbf{B})_1 + z(\hat{\mathbf{r}} \times \mathbf{C})_1 \right) \dot{V}_1 - (\mathbf{a} \times \boldsymbol{\alpha})_1 + b_0 \nabla \cdot \mathbf{A}, \end{aligned} \quad (4.30)$$

$$\begin{aligned} \Phi_{0y} = & \left((yB_{3x} - xB_{3y}) + (xB_{2z} - zB_{2x}) + (zB_{1y} - yB_{1z}) + (A_3 - C_1) \right) V_1 \\ & - \left(x(\hat{\mathbf{r}} \times \mathbf{A})_2 + y(\hat{\mathbf{r}} \times \mathbf{B})_2 + z(\hat{\mathbf{r}} \times \mathbf{C})_2 \right) \dot{V}_1 - (\mathbf{a} \times \boldsymbol{\alpha})_2 + b_0 \nabla \cdot \mathbf{B}, \end{aligned} \quad (4.31)$$

$$\begin{aligned} \Phi_{0z} = & \left((yC_{3x} - xC_{3y}) + (xC_{2z} - zC_{2x}) + (zC_{1y} - yC_{1z}) + (B_1 - A_2) \right) V_1 \\ & - \left(x(\hat{\mathbf{r}} \times \mathbf{A})_3 + y(\hat{\mathbf{r}} \times \mathbf{B})_3 + z(\hat{\mathbf{r}} \times \mathbf{C})_3 \right) \dot{V}_1 - (\mathbf{a} \times \boldsymbol{\alpha})_3 + b_0 \nabla \cdot \mathbf{C}, \end{aligned} \quad (4.32)$$

where $\dot{V}_1 \equiv dV_1/dr$, $\hat{\mathbf{r}}$ is the unit displacement vector, $\mathbf{a} = (a_1, a_2, a_3)$ and $\mathbf{b} = (b_1, b_2, b_3)$ are the space parts of the Lorentz violating parameters a_μ and b_μ respectively, and $\mathbf{A} = (A_1, A_2, A_3)$ (\mathbf{B} and \mathbf{C} are defined similarly) is introduced in the integral of motion X . The rest of the determining equations from the coefficients of the first-order terms can be expressed in terms of certain derivative combinations of the determining equations from the coefficients of the second-order terms given in eqs. (4.3)–(4.20). For completeness they are symbolically listed below:

$$2(\text{Eq}(4.9))_x - (\text{Eq}(4.17))_y - (\text{Eq}(4.19))_z = 0, \quad (4.33)$$

$$2(\text{Eq}(4.4))_x - (\text{Eq}(4.18))_y + (\text{Eq}(4.13))_z = 0, \quad (4.34)$$

$$2(\text{Eq}(4.3))_x + (\text{Eq}(4.12))_y - (\text{Eq}(4.15))_z = 0, \quad (4.35)$$

$$2(\text{Eq}(4.5))_y + (\text{Eq}(4.17))_x + (\text{Eq}(4.14))_z = 0, \quad (4.36)$$

$$2(\text{Eq}(4.10))_y + (\text{Eq}(4.18))_x - (\text{Eq}(4.20))_z = 0, \quad (4.37)$$

$$2(\text{Eq}(4.6))_y + (\text{Eq}(4.12))_x - (\text{Eq}(4.16))_z = 0, \quad (4.38)$$

$$2(\text{Eq}(4.7))_z + (\text{Eq}(4.19))_x + (\text{Eq}(4.14))_y = 0, \quad (4.39)$$

$$2(\text{Eq}(4.8))_z + (\text{Eq}(4.13))_x + (\text{Eq}(4.20))_y = 0, \quad (4.40)$$

$$2(\text{Eq}(4.11))_z + (\text{Eq}(4.15))_x + (\text{Eq}(4.16))_y = 0, \quad (4.41)$$

where $(\dots)_x$ for example represents the derivative of the entire equation with respect to x .

We do not include the expressions for the 8 determining differential equations coming from the zeroth-order terms. They are very lengthy and their exact form is not particularly illuminating. We simplify the equations by imposing the solutions obtained from the second- and first-order terms, and then they are either trivially satisfied or reduce to simple constraint equations (like eq. (3.7) in the 2D case), presented for each case in the following discussion.

In all three cases to be discussed below we have the same integral of motion eq. (2.6) and the functions A_0 , B_0 and C_0 are obtained the same as in eq. (4.2). Therefore, we will briefly discuss the solutions for the rest of the coefficients by applying the technique used so far.

4.1 Lorentz violation with vector coupling ($a_i \neq 0$):

In this case we have the Hamiltonian eq. (2.5) with $\mathbf{b} = \mathbf{0}$ and $b_0 = 0$. It is found by using Maple that when $b_0 = 0$ we have the following solution for the 18 determining equations given in eq. (4.3)–(4.20)

$$\begin{aligned} A_1 &= 0, & A_2 &= zw, & A_3 &= -yw, \\ B_1 &= -zw, & B_2 &= 0, & B_3 &= xw, \\ C_1 &= yw, & C_2 &= -xw, & C_3 &= 0, \end{aligned} \tag{4.42}$$

where w is an integration constant.

After introducing eq. (4.42) into the rest of the determining equations it is immediately found from eq. (4.30)–(4.32) that

$$\Phi_0 = (\alpha_2 a_3 - \alpha_3 a_2)x + (\alpha_3 a_1 - \alpha_1 a_3)y + (\alpha_1 a_2 - \alpha_2 a_1)z, \tag{4.43}$$

and then we are left with 9 first-order partial differential equations for $\Phi = (\Phi_1, \Phi_2, \Phi_3)$. These are the equations eq. (4.21)–(4.29) with $\mathbf{b} = \mathbf{0}$ and $b_0 = 0$. In order to solve this system we express the first-derivatives of Φ and look for the compatibility of the mixed partial derivatives. The requirement of the equality of the mixed partial derivatives gives another 9 equations for Φ and its first-order derivatives. Now, introducing the first-derivatives of Φ which are found from eq. (4.21)–(4.29) into this system, we get a system of algebraic equations for Φ . In order to have a solution of this algebraic system we must have $\beta = \mathbf{0}$ and $w = 0$. It is not surprising to require $w = 0$ since it is the coefficient of the $\sigma \cdot \mathbf{L}$ term in eq. (2.6) and clearly it cannot commute with the Hamiltonian eq. (2.5) in the presence of \mathbf{a} terms. Setting $\beta = \mathbf{0}$ and $w = 0$ we find

$$\Phi = \frac{\alpha}{2}. \tag{4.44}$$

Finally, if we introduce all the information gathered from the coefficients of the higher-order terms into the determining equations coming from the coefficients of the zeroth-order terms, we find that the following 3 conditions must also be satisfied

$$\begin{aligned} a_1(y\alpha_2 + z\alpha_3) - (a_2y + a_3z)\alpha_1 &= 0, \\ a_2(x\alpha_1 + z\alpha_3) - (a_1x + a_3z)\alpha_2 &= 0, \\ a_3(x\alpha_1 + y\alpha_2) - (a_1x + a_2y)\alpha_3 &= 0. \end{aligned} \tag{4.45}$$

Clearly it is seen that in order to satisfy these 3 conditions we must either have $\mathbf{a} = \mathbf{0}$ or $\boldsymbol{\alpha} = \mathbf{0}$. Since the former case implies that there are no LV terms and the latter one indicates we have no nontrivial first-order integral of motion, we conclude that in the presence of \mathbf{a} terms there is no first-order integrability unlike the 2D case, if \mathbf{a} is perpendicular to the plane of the motion (L_3 as discussed is constant of motion). It should be noted that a_μ term can be eliminated with a transformation and in general is not observable physically. This is only true for simple one fermion models or non-interacting multi-fermion cases but does not hold for a generic multi-fermion theory.

4.2 Lorentz violation with pure spacelike axial-vector coupling ($b_i \neq 0$):

After setting $\mathbf{a} = \mathbf{0}$ and $b_0 = 0$ in eq. (2.5), we again find the same solution as in eq. (4.42) for \mathbf{A}, \mathbf{B} , and \mathbf{C} since the 18 equations are identical in the absence of b_0 . However, from eq. (4.30)–(4.32), in this case we obtain $\Phi_0 = \text{const}$, which can be taken as zero without loss of generality. Then, we proceed in a similar fashion as in the previous case and reach the same conclusion for the system of 9 differential equations given in eq. (4.21)–(4.29), with $\mathbf{a} = \mathbf{0}$ and $b_0 = 0$. That is, $\boldsymbol{\Phi} = \boldsymbol{\alpha}/2$ is still the solution together with the requirement that $\boldsymbol{\beta} = \mathbf{0}$ and $w = 0$. After introducing all the information gathered from the coefficients of the higher-order terms into the determining equations coming from the coefficients of the zeroth-order terms, we find, as in the previous case, that the following condition must also be satisfied

$$\mathbf{b} \cdot \boldsymbol{\alpha} = 0. \tag{4.46}$$

Again, we can conclude that in general in the presence of \mathbf{b} terms there is no first-order integrability. However, we may find some integral of motions in some special cases (e.g. choosing $b_1 = 0, b_2 = 0$ and $\alpha_1 = 0, \alpha_2 = 0$). In general there is a constant of motion $X_{3D}^{(b_i \neq 0)} = \boldsymbol{\alpha} \cdot \mathbf{J}$ if \mathbf{b} is a vector perpendicular to $\boldsymbol{\alpha}$ (here \mathbf{J} represents the total angular momentum operator). Of course, not all components of $\boldsymbol{\alpha}$ are independent due to eq. (4.46). For example, if we assume the i^{th} component of \mathbf{b} to be nonzero, then one can alternatively write the constant of motion as

$$X_{3D}^{(b_i \neq 0)} = \frac{1}{b_i} [(\boldsymbol{\alpha} \times (\mathbf{b} \times \mathbf{J}))_i]. \tag{4.47}$$

where $\boldsymbol{\alpha}$ is otherwise arbitrary. From here one could think that there are two constants of motion after eliminating one of the components of $\boldsymbol{\alpha}$ but this is not true. Once the orientation of \mathbf{b} is fixed, there is a unique $\boldsymbol{\alpha}$ so that we have one additional constant of motion, which could be chosen as J_3 . One noticeable difference from the 2D case is that the total angular momentum J_3 is now conserved, but this is not enough to make the system first-order integrable.

4.3 Lorentz violation with pure timelike axial-vector coupling ($b_0 \neq 0$):

In this case, the set of 18 equations in eqs. (4.3)–(4.20) differs from the previous 2 cases since $b_0 \neq 0$. The only solution of this system is the so called *trivial solution* which is $\mathbf{A} = \mathbf{0}, \mathbf{B} = \mathbf{0}$ and $\mathbf{C} = \mathbf{0}$. We find also that $\Phi_0 = \text{const}$, which can be taken as zero

without loss of generality, and proceed in the same way as in the previous cases. We again find the same Φ as in eq. (4.44) together with the only requirement $\beta = \mathbf{0}$. However, this time the determining equations coming from the coefficients of the zeroth-order terms are identically satisfied if we introduce all the information gathered from the coefficients of the higher-order terms. The form of $X_{3D}^{(b_0 \neq 0)}$ is the same as in the previous case but without the constraint eq. (4.46). There exist however two additional constants of motion since the third one is related to the first two through an $SU(2)$ algebra.

Thus, we conclude that the Hamiltonian eq. (2.5) with $\mathbf{a} = \mathbf{0}$ and $\mathbf{b} = \mathbf{0}$ is first-order integrable even though it has LV terms ($b_0 \neq 0$). It is easily seen that the components of \mathbf{J} commute with H , hence the system is integrable. We may choose $\{J_i, \mathbf{J}^2, H\}$ as a commuting set of operators. It is important to note that in the commuting set we choose \mathbf{J}^2 but there is no way to get it in $X_{3D}^{(b_0 \neq 0)}$ since \mathbf{J}^2 is a second-order operator. However, first-order integrals of motion J_i generate \mathbf{J}^2 .

5. Perturbative solutions

The whole idea in this study is to seek, with a systematic method, the first-order integrability/superintegrability of the Pauli system in (axial-)vectorial LV background. The ultimate goal is to achieve at least a quasi-exact solution. This could be important for the $b_0 \neq 0$ case since the bound on b_0 for electron is much weaker than both \mathbf{a} and \mathbf{b} , and if the effective size of the system comparable with b_0^{-1} , the perturbative approach fails. So, it would be enough to mainly concentrate on the $b_0 \neq 0$ case and even perform a second-order analysis. However, our experience is that it would be much harder to solve the equations of overdetermined systems analytically, especially if one keeps the spin-orbit term.

Based on our analysis we don't have first-order superintegrability in either 2D or 3D. There are first-order integrable cases in 2D under special arrangements but this is only true in 3D for the $b_0 \neq 0$ case, where exact solutions could be of interest. Of course, neither integrability nor superintegrability does guarantee the separability and existence of exact solution but either is an important step in finding one. Recently, a perturbative approach has been carried out for a similar Hamiltonian system in the same LV background within the QED version of Standard Model Extension [18, 19]. For completeness, we like to briefly summarize their results. Note that the analysis in ref. [18] includes an external magnetic field but not the spin-orbit term. A more extensive perturbative analysis for all types of Lorentz violating couplings is found in [20]. There are also studies [21] on the violation of Einstein Equivalence Principle, which in principle may lead to Lorentz violation, by considering some non-gravitational interactions such as Lamb shift and the anomalous magnetic moment $g - 2$ of electron in non-metric theories.

For the case with a_μ , as discussed in section 4.1, using a field redefinition by a factor of $\exp(i a_\mu x^\mu)$, the energy is shifted by a constant a_0 and wavefunction has an overall phase factor $\exp(i \mathbf{a} \cdot \mathbf{r})$, compared to a hydrogen atom without LV background.

For the case $\mathbf{b} \neq 0$, the perturbative approach is applied [18] and energy shifts are obtained, by an amount proportional to $b_3 m_s$ (to first order). Here m_s is the magnetic spin quantum number. The wave function is unchanged. The energy correction is proportional

to $b_3 m_j / (l + 1)$ if the spin-orbit correction is included. Here m_j and l are the total magnetic angular momentum and the orbital angular momentum numbers, respectively. These corrections are quite small due to experimental bounds on $|\mathbf{b}|$.

The case with $b_0 \neq 0$ is indeed interesting. It is shown that using a unitarity transformation one could relate a real hydrogen with all corrections (spin-orbit, relativistic, and the so-called Darwin term) to the one with additional LV background of the form $b_0 \boldsymbol{\sigma} \cdot \mathbf{p}$. The unitarity transformation is $U = 1 - i b_0 (1 + \Sigma/r) \boldsymbol{\sigma} \cdot \mathbf{r}$, up to first order in b_0 , so that $H_{\text{real Hy}}^{(b_0 \neq 0)} = U^\dagger H_{\text{real Hy}} U$ and $\Psi^{(b_0 \neq 0)} = U^\dagger \Psi$. Here Σ is a constant appropriately chosen to obtain the correction terms. It is then straightforward to solve the system $H_{\text{real Hy}}$ by applying perturbation theory and to transform everything back to the original system. As usual there is no contribution to the energy spectrum of the hydrogen atom from the b_0 term, up to first order in b_0 . However, the wave function develops a b_0 dependent part. See the ref. [18] for the details.

6. Conclusion

In this study we have analyzed in a systematic way the integrability of a Pauli system in a LV background both in two- and three-dimensions. We consider the Hamiltonian which originates from the non-relativistic limit of the QED part of the so-called Standard Model Extension. We kept only two types of LV couplings to the electron, the vector type of coupling represented by a_μ and the axial-vector type of coupling by b_μ . Since a_0 enters as a constant term in the Hamiltonian, we consider in both 2D and 3D the following three separate cases; LV from $\mathbf{a} \neq 0$, LV from $\mathbf{b} \neq 0$, and LV from $b_0 \neq 0$. The last case is especially important since perturbative approaches could fail under certain circumstances, due to weak experimental bounds on b_0 , and exact solutions might be required. Note that in our study we kept the spin-orbit correction to the hydrogen atom, partially because the correction from the LV term with $b_0 \neq 0$ could be comparable with the spin-orbit term, as shown in section 2.

After writing the most general constant of motion operator X in first-order derivatives, we obtained sets of overdetermined differential equations from the vanishing of the commutator of X with the Hamiltonian. In 2D, first-order integrability is possible if \mathbf{a} is perpendicular to the plane of the motion, when the LV is due to $\mathbf{a} \neq 0$ only. The constant of motion is one of the components of the angular momentum operator \mathbf{L} (L_3). Our conclusion is the same for $\mathbf{b} \neq 0$ case, with the difference that the vector \mathbf{b} can have arbitrary components in each direction. The last case in 2D is $b_0 \neq 0$ and first-order integrability is restored like in the other cases with one difference: the z -component of the total angular momentum operator J_3 is conserved if the motion is in xy plane.

Based on our 2D results, we extended our discussion into a more realistic 3D picture by following the same procedure. The first-order integrability disappears for the $\mathbf{a} \neq 0$ case in 3D. For $\mathbf{b} \neq 0$ case, even though J_3 becomes a constant of motion, it is not enough to make the system integrable (since we need another one, unlike 2D case). The first-order integrability in 3D is retained only for $\mathbf{a} = \mathbf{b} = 0$ with $b_0 \neq 0$ where we can find two additional constant of motion operators like J_1 and J_2 . We believe that studying the

$b_0 \neq 0$ case up to second-order is worthwhile in search of superintegrability. This is in fact the case where exact solutions could be physically relevant. One caveat is that, at the second order, solving the overdetermined partial differential equations becomes much more challenging. For completeness, we also summarized the results of a recent paper [18] where unitary transformations and perturbative approaches are employed for similar systems.

Acknowledgments

We thank Alan Kostelecky and Pavel Winternitz for communications and helpful discussions. The work of M.F. and I.T. is supported in part by NSERC of Canada under the Grant No. SAP01105354. Í.Y. acknowledges a postdoctoral fellowship awarded by the Laboratory of Mathematical Physics of the CRM, Université de Montréal.

References

- [1] V. Bargmann, *Theory of the hydrogen atom*, *Z. Phys.* **99** (1936) 576;
V.A. Fock, *To the theory of the hydrogen atom*, *Z. Phys.* **98** (1935) 145.
- [2] J. Jauch and E. Hill, *On the problem of degeneracy in quantum mechanics*, *Phys. Rev.* **57** (1940) 641;
M. Moshinsky and Y.F. Smirnov, *The harmonic oscillator in modern physics*, Harwood, Amsterdam (1996).
- [3] I. Friš, V. Mandrosov, J. Smorodinsky, M. Uhlř and P. Winternitz, *On higher-order symmetries in quantum mechanics*, *Phys. Lett.* **16** (1965) 354;
P. Winternitz, J. Smorodinsky, M. Uhlř and I. Friš, *Symmetry groups in classical and quantum mechanics*, *Yad. Fiz.* **4** (1966) 625 [*Sov. J. Nucl. Phys.* **4** (1967) 444];
A. Makarov, J. Smorodinsky, K. Valiev and P. Winternitz, *A systematic search for non-relativistic system with dynamical symmetries*, *Nuovo Cim.* **A 52** (1967) 1061;
N.W. Evans, *Superintegrability in classical mechanics*, *Phys. Rev.* **A 41** (1990) 5666;
Superintegrability of the Winternitz system, *Phys. Lett.* **A 147** (1990) 483.
- [4] E.G. Kalnins, J.M. Kress and J.W. Miller, *Second-order superintegrable systems in conformally flat spaces. I. Two-dimensional classical structure theory*, *J. Math. Phys.* **46** (2005) 053509; *Second order superintegrable systems in conformally flat spaces. II. The classical two-dimensional Stäckel transform*, *J. Math. Phys.* **46** (2005) 053510; *Second order superintegrable systems in conformally flat spaces. III. Three-dimensional classical structure theory*, *J. Math. Phys.* **46** (2005) 103507;
M.B. Sheftel, P. Tempesta and P. Winternitz, *Superintegrable systems in quantum mechanics and classical Lie theory*, *J. Math. Phys.* **42** (2001) 659;
P. Tempesta, A.V. Turbiner and P. Winternitz, *Exact solvability of superintegrable systems*, *J. Math. Phys.* **42** (2001) 4248 [[hep-th/0011209](#)].
- [5] B. Dorizzi, B. Grammaticos, A. Ramani and P. Winternitz, *Integrable Hamiltonian systems with velocity dependent potentials*, *J. Math. Phys.* **26** (1985) 3070;
J. Bérubé and P. Winternitz, *Integrable and superintegrable quantum systems in a magnetic field*, *J. Math. Phys.* **45** (2004) 1959;
G. Pucacco and K. Rosquist, *Integrable Hamiltonian systems with vector potentials*, *J. Math. Phys.* **46** (2005) 012701;

- P. Tempesta et al., *Superintegrability in classical and quantum systems*, CRM Proc. Lecture Notes **37**, Providence U.S.A. (2004).
- [6] P. Winternitz and İ. Yurduşen, *Integrable and superintegrable systems with spin*, *J. Math. Phys.* **47** (2006) 103509 [[math-ph/0604050](#)]; *Superintegrable systems with spin in two- and three-dimensional Euclidean spaces*, in G. Gaeta, R. Vitolo and S. Walcher, *Symmetry and perturbation theory: proceedings of the international conference on SPT 2007*, World Scientific, Singapore (2007) [[arXiv:0711.0753](#)].
- [7] D. Colladay and V.A. Kostelecky, *CPT violation and the standard model*, *Phys. Rev.* **D 55** (1997) 6760 [[hep-ph/9703464](#)]; *Lorentz-violating extension of the standard model*, *Phys. Rev.* **D 58** (1998) 116002 [[hep-ph/9809521](#)].
- [8] V.A. Kostelecky, *Gravity, Lorentz violation and the standard model*, *Phys. Rev.* **D 69** (2004) 105009 [[hep-th/0312310](#)];
R. Bluhm and V.A. Kostelecky, *Spontaneous Lorentz violation, Nambu-Goldstone modes and gravity*, *Phys. Rev.* **D 71** (2005) 065008 [[hep-th/0412320](#)];
Q.G. Bailey and V.A. Kostelecky, *Signals for Lorentz violation in post-newtonian gravity*, *Phys. Rev.* **D 74** (2006) 045001 [[gr-qc/0603030](#)];
V.A. Kostelecky and R. Potting, *Gravity from local Lorentz violation*, *Gen. Rel. Grav.* **37** (2005) 1675 [*Int. J. Mod. Phys.* **D 14** (2005) 2341] [[gr-qc/0510124](#)].
- [9] R. Bluhm, V.A. Kostelecky and N. Russell, *CPT and Lorentz tests in hydrogen and antihydrogen*, *Phys. Rev. Lett.* **82** (1999) 2254 [[hep-ph/9810269](#)]; *Hydrogen and antihydrogen spectroscopy for studies of CPT and Lorentz symmetry*, [hep-ph/9810327](#); *Searching for Lorentz violation in the ground state of hydrogen*, [hep-ph/0003223](#);
R. Bluhm, *Lorentz and CPT tests in atomic systems*, *AIP Conf. Proc.* **539** (2000) 109 [[hep-ph/0006033](#)]. R. Bluhm, V.A. Kostelecky and N. Russell, *Searching for Lorentz violation in the ground state of hydrogen*, [hep-ph/0003223](#).
- [10] V.A. Kostelecky, *CPT and Lorentz Symmetry*, World Scientific, Singapore (1999); *CPT and Lorentz Symmetry II*, World Scientific, Singapore (2002); *CPT and Lorentz Symmetry III*, World Scientific, Singapore (2005).
- [11] R. Bluhm, *Overview of the SME: implications and phenomenology of Lorentz violation*, *Lect. Notes Phys.* **702** (2006) 191 [[hep-ph/0506054](#)];
D. Mattingly, *Modern tests of Lorentz invariance*, *Living Rev. Rel.* **8** (2005) 5 [[gr-qc/0502097](#)];
D.L. Anderson, M. Sher and I. Turan, *Lorentz and CPT violation in the Higgs sector*, *Phys. Rev.* **D 70** (2004) 016001 [[hep-ph/0403116](#)];
G. Amelino-Camelia, C. Lammerzahl, A. Macias and H. Muller, *The search for quantum gravity signals*, *AIP Conf. Proc.* **758** (2005) 30 [[gr-qc/0501053](#)];
H. Vucetich, *Testing Lorentz invariance violation in quantum gravity theories*, [gr-qc/0502093](#).
- [12] V.A. Kostelecky and C.D. Lane, *Nonrelativistic quantum hamiltonian for Lorentz violation*, *J. Math. Phys.* **40** (1999) 6245 [[hep-ph/9909542](#)];
R. Lehnert, *Dirac theory within the standard-model extension*, *J. Math. Phys.* **45** (2004) 3399 [[hep-ph/0401084](#)];
Q.G. Bailey and V.A. Kostelecky, *Lorentz-violating electrostatics and magnetostatics*, *Phys. Rev.* **D 70** (2004) 076006 [[hep-ph/0407252](#)];
A. Kostelecky and M. Mewes, *Lorentz-violating electrodynamics and the cosmic microwave background*, *Phys. Rev. Lett.* **99** (2007) 011601 [[astro-ph/0702379](#)].

- [13] R. Jackiw and V.A. Kostelecky, *Radiatively induced Lorentz and CPT violation in electrodynamics*, *Phys. Rev. Lett.* **82** (1999) 3572 [[hep-ph/9901358](#)];
M. Pérez-Victoria, *Physical (ir)relevance of ambiguities to Lorentz and CPT violation in QED*, *JHEP* **04** (2001) 032 [[hep-th/0102021](#)];
V.A. Kostelecky, C.D. Lane and A.G.M. Pickering, *One-loop renormalization of Lorentz-violating electrodynamics*, *Phys. Rev.* **D 65** (2002) 056006 [[hep-th/0111123](#)];
B. Altschul, *Failure of gauge invariance in the nonperturbative formulation of massless Lorentz-violating QED*, *Phys. Rev.* **D 69** (2004) 125009 [[hep-th/0311200](#)]; *Gauge invariance and the Pauli-Villars regulator in Lorentz- and CPT-violating electrodynamics*, *Phys. Rev.* **D 70** (2004) 101701 [[hep-th/0407172](#)];
V.A. Kostelecky, R. Lehnert and M.J. Perry, *Spacetime-varying couplings and Lorentz violation*, *Phys. Rev.* **D 68** (2003) 123511 [[astro-ph/0212003](#)];
H. Muller, C. Braxmaier, S. Herrmann, A. Peters and C. Laemmerzahl, *Electromagnetic cavities and Lorentz invariance violation*, *Phys. Rev.* **D 67** (2003) 056006 [[hep-ph/0212289](#)];
M. Frank and I. Turan, *The Casimir force in a Lorentz violating theory*, *Phys. Rev.* **D 74** (2006) 033016 [[hep-ph/0607087](#)];
H. Belich, T. Costa-Soares, M.M. Ferreira and J.A. Helayel-Neto, *Classical solutions in a Lorentz-violating scenario of Maxwell-Chern-Simons-Proca electrodynamics*, *Eur. Phys. J.* **C 42** (2005) 127 [[hep-th/0411151](#)];
T. Mariz, J.R. Nascimento, E. Passos, R.F. Ribeiro and F.A. Brito, *A remark on Lorentz violation at finite temperature*, *JHEP* **10** (2005) 019 [[hep-th/0509008](#)].
- [14] C. Adam and F.R. Klinkhamer, *Photon decay in a CPT-violating extension of quantum electrodynamics*, *Nucl. Phys.* **B 657** (2003) 214 [[hep-th/0212028](#)];
T. Jacobson, S. Liberati and D. Mattingly, *Threshold effects and Planck scale Lorentz violation: combined constraints from high energy astrophysics*, *Phys. Rev.* **D 67** (2003) 124011 [[hep-ph/0209264](#)];
V.A. Kostelecky and A.G.M. Pickering, *Vacuum photon splitting in Lorentz-violating quantum electrodynamics*, *Phys. Rev. Lett.* **91** (2003) 031801 [[hep-ph/0212382](#)];
R. Lehnert and R. Potting, *Vacuum Cerenkov radiation*, *Phys. Rev. Lett.* **93** (2004) 110402 [[hep-ph/0406128](#)]; *The Cerenkov effect in Lorentz-violating vacua*, *Phys. Rev.* **D 70** (2004) 125010 [*Erratum ibid.* **D 70** (2004) 129906] [[hep-ph/0408285](#)];
T.A. Jacobson, S. Liberati, D. Mattingly and F.W. Stecker, *New limits on Planck scale Lorentz violation in QED*, *Phys. Rev. Lett.* **93** (2004) 021101 [[astro-ph/0309681](#)];
B. Altschul, *Lorentz violation and synchrotron radiation*, *Phys. Rev.* **D 72** (2005) 085003 [[hep-th/0507258](#)];
F.R. Klinkhamer and C. Rupp, *Photon-propagation model with random background field: length scales and Cherenkov limits*, *Phys. Rev.* **D 72** (2005) 017901 [[hep-ph/0506071](#)];
C. Kaufhold and F.R. Klinkhamer, *Vacuum Cherenkov radiation and photon triple-splitting in a Lorentz-noninvariant extension of quantum electrodynamics*, *Nucl. Phys.* **B 734** (2006) 1 [[hep-th/0508074](#)].
- [15] O.W. Greenberg, *CPT violation implies violation of Lorentz invariance*, *Phys. Rev. Lett.* **89** (2002) 231602 [[hep-ph/0201258](#)]; *Hybrid Dirac fields*, *Phys. Lett.* **B 567** (2003) 179 [[hep-ph/0305276](#)].
- [16] D. Colladay and P. McDonald, *Redefining spinors in Lorentz-violating QED*, *J. Math. Phys.* **43** (2002) 3554 [[hep-ph/0202066](#)].
- [17] K.M. Heal, M. Hansen and K. Rickard, *Maple V learning guide for release 5*, Springer Verlag (1997);

see also references from <http://www.maplesoft.com/maplebooks.htm>.

- [18] O.G. Kharlanov and V.C. Zhukovsky, *CPT and Lorentz violation effects in hydrogen-like atoms*, [arXiv:0705.3306](#).
- [19] M.M. Ferreira and F.M.O. Moucherek, *Influence of Lorentz- and CPT-violating terms on the Dirac equation*, *Int. J. Mod. Phys. A* **21** (2006) 6211 [[hep-th/0601018](#)];
M.M. Ferreira, A.R. Gomes and R.C.C. Lopes, *Influence of Lorentz-violating terms on a two-level system*, *Phys. Rev. D* **76** (2007) 105031 [[arXiv:0707.4660](#)];
H. Belich, T. Costa-Soares, J. Ferreira, M. M., J.A. Helayel-Neto and F.M.O. Mouchereck, *Lorentz-violating corrections on the hydrogen spectrum induced by a non-minimal coupling*, *Phys. Rev. D* **74** (2006) 065009 [[hep-th/0604149](#)].
- [20] R. Bluhm, V.A. Kostelecky and N. Russell, *CPT and Lorentz tests in penning traps*, *Phys. Rev. D* **57** (1998) 3932 [[hep-ph/9809543](#)].
- [21] C. Alvarez and R.B. Mann, *Testing the equivalence principle by lamb shift energies*, *Phys. Rev. D* **54** (1996) 5954 [[gr-qc/9507040](#)];
The equivalence principle and anomalous magnetic moment experiments, *Phys. Rev. D* **54** (1996) 7097 [[gr-qc/9511028](#)];
The equivalence principle and g-2 experiments, *Phys. Lett. B* **409** (1997) 83 [[gr-qc/9510070](#)].