

Title: Effects of water temperature on summer periphyton biomass in shallow lakes: a pan-European mesocosm experiment

Author(s): Mahdy, A., Hilt, S., Filiz, N., Beklioğlu, M., Hejzlar, J., Özkundakci, D., ... Adrian, R.

Document type: Postprint

Terms of Use: Copyright applies. A non-exclusive, non-transferable and limited right to use is granted. This document is intended solely for personal, non-commercial use.

Citation: Mahdy, A., Hilt, S., Filiz, N., Beklioğlu, M., Hejzlar, J., Özkundakci, D., ... Adrian, R. (2015). Effects of water temperature on summer periphyton biomass in shallow lakes: a pan-European mesocosm experiment. *Aquatic Sciences*, 77(3), 499–510.
<https://doi.org/10.1007/s00027-015-0394-7>

This is a post-peer-review, pre-copyedit version of an article published in *Aquatic Sciences*. The final authenticated version is available online at:
<http://dx.doi.org/10.1007/s00027-015-0394-7>

1 **Effects of water temperature on summer periphyton biomass in shallow lakes:**
2 **a pan-European mesocosm experiment**

3
4 Aldoushy Mahdy^{a,b,c}, Sabine Hilt^{a*}, Nur Filiz^d, Meryem Beklioğlu^d, Josef Hejzlar^e, Deniz
5 Özkundakci^a, Eva Papastergiadou^f, Ulrike Scharfenberger^a, Michal Šorf^{e,g}, Kostas Stefanidis^f,
6 Lea Tuvikene^h, Priit Zingel^h, Martin Søndergaardⁱ, Erik Jeppesen^{i,j}, Rita Adrian^{a,b}

7
8 ^aLeibniz Institute of Freshwater Ecology and Inland Fisheries, IGB, Berlin, Germany

9 ^bFree University Berlin, Department of Biology, Chemistry and Pharmacy, Berlin, Germany

10 ^cDepartment of Zoology, Faculty of Science, Al-Azhar University (Assiut Branch), Assiut
11 71524, Egypt

12 ^dMiddle East Technical University, Biological Sciences Department, Limnology Laboratory,
13 Ankara, Turkey

14 ^eBiology Centre of the Academy of Sciences of the Czech Republic, Institute of
15 Hydrobiology, České Budějovice, Czech Republic

16 ^fUniversity of Patras, Department of Biology, Patras, Greece

17 ^gFaculty of Science, University of South Bohemia, České Budějovice, Czech Republic

18 ^hCentre for Limnology, Estonian University of Life Sciences, 61117 Rannu, Tartu County,
19 Estonia

20 ⁱDepartment of Bioscience, Aarhus University, Vejløvej 25, 8600 Silkeborg, Denmark

21 ^jSino-Danish Centre for Education and Research, Beijing, China

22
23 *corresponding author: hilt@igb-berlin.de, tel.: +49 30 64181677, fax: +49 30 64181682

24 **Abstract**

25 Periphyton communities play an important role in shallow lakes and are controlled by direct
26 forces such as temperature, light, nutrients, and invertebrate grazing, but also indirectly by
27 planktivorous fish predation. We performed a pan-European lake mesocosm experiment on
28 periphyton colonization covering five countries along a north/south geographical/temperature
29 gradient (Estonia, Germany, Czech Republic, Turkey, and Greece). Periphyton biomass on
30 artificial polypropylene strips exposed at 50 cm water depth at low and high nutrient regimes
31 (with mean total phosphorus concentration of 20 and 65 $\mu\text{g L}^{-1}$, respectively) was compared
32 during mid-summer. No significant effect of nutrient loading on periphyton biomass was
33 observed as nutrient concentrations in the mesocosms were generally above limiting values.
34 Water temperature significantly enhanced summer periphyton biomass development.
35 Additionally, direct and indirect top-down control of snails and fish emerged as a significant
36 factor in periphyton biomass control.

37

38 **Keywords:** climate change, epiphyton, eutrophication, grazing, top-down control

39 **Introduction**

40 Shallow lakes tend to exist in one of two stable states, a macrophyte-dominated state with
41 high water transparency or a turbid, phytoplankton-dominated state without submerged
42 macrophytes (Scheffer et al. 1993). Phillips et al. (1978) found that lakes that had switched
43 from a macrophyte-dominated clear state to a phytoplankton-dominated turbid state during a
44 period of eutrophication showed an increase in periphyton biomass prior to phytoplankton
45 development. Jones and Sayer (2003) supported these findings of increased periphyton
46 shading as the first step leading to the decline of submerged macrophytes in eutrophic lakes.
47 However, they suggested that a top-down control cascade from fish via scraping invertebrates
48 rather than nutrient concentrations would be responsible for periphyton control under
49 eutrophic conditions. Liboriussen et al. (2005) also found a significant top-down control of
50 periphyton biomass in a mesocosm experiment in Denmark. The relative importance of
51 bottom-up and top-down control of periphyton biomass may, though, vary widely across
52 spatial (i.e. between lakes) and temporal (i.e. between years) scales (Jeppesen et al. 1997).

53 Climate regimes are likely to affect lake biota communities (IPCC 2013). In
54 productive lakes, climate warming accelerates a shift in trophic state (Mooij et al. 2005;
55 Adrian et al. 2009) and consequently affects light conditions. The subsequent responses by
56 plankton communities have been studied extensively (Adrian et al. 2006; Seebens et al. 2009;
57 Wagner and Adrian 2009). How periphyton growth is affected by warming is debated and the
58 results obtained so far are ambiguous. Some studies have shown an increase in periphyton
59 biomass with increasing water temperature (Tarkowska-Kukuryk and Mieczan 2012; Patrick
60 et al. 2012), while Shurin et al. (2012) in a mesocosm study demonstrated that periphyton
61 chlorophyll *a* declined with elevated temperatures (3 °C above ambient). Such differences
62 may be attributed to variations in the grazing pressure by invertebrates and fish. In microcosm
63 experiments, Cao et al. (2014) observed as response to increased temperatures an increase in
64 periphyton biomass when snails were absent but no effect when snails were present. McKee

65 et al. (2002) also found grazers to benefit more than periphyton from enhanced temperatures.
66 Moreover, herbivory and omnivory among fishes increase with temperature (González-
67 Bergonzoni et al. 2012; Meerhoff et al. 2012), and many fish species (or size classes of fish)
68 feed on periphyton in subtropical and tropical lakes (Teixeira-de Mello et al. 2009).

69 Comparative studies on periphyton dynamics along latitudinal scales are scarce.
70 Bécares et al. (2008) conducted a mesocosm experiment across a European latitudinal
71 gradient from Finland to Spain and found that periphyton chlorophyll *a* concentrations were
72 overall positively related to nutrient loading. Top-down effects by fish were significant only
73 in a few sites and were assumed to be related to their contribution to the nutrient pool. Under
74 these conditions southern lakes exhibited lower periphyton densities than northern lakes
75 because of the larger phytoplankton biomass in the south and its shading effects on periphyton
76 at similar nutrient loadings. In a comparative experimental field study by Meerhoff et al.
77 (2007), a substantially lower periphyton biomass on artificial plants was found in lakes in
78 subtropical Uruguay than in temperate Denmark. Despite a much lower biomass of
79 invertebrate periphyton grazers due to high fish predation in Uruguay, the authors attributed
80 the lower periphyton biomass in the warm lakes to direct control by fish grazing. Therefore,
81 periphyton biomass might be directly or indirectly affected by nutrients, temperature, grazers,
82 and fish, but the mechanisms of the underlying processes and potential interactions are still
83 poorly understood.

84 We studied periphyton development on artificial polypropylene strips exposed in
85 mesocosms with two different nutrient loadings resembling mesotrophic and eutrophic
86 conditions and at moderate fish density in five European countries (Estonia, Germany, Czech
87 Republic, Turkey, and Greece). A latitudinal temperature gradient was expected and effects
88 on periphyton biomass were studied for a period of one month in July and August 2011. We
89 hypothesize that higher nutrient loading and warmer temperatures increase summer
90 periphyton biomass under these conditions.

91 **Materials and methods**

92 Study sites and experimental set-up

93 We conducted a mesocosm experiment in five countries across Europe covering a climate
94 gradient from Estonia (58° N 26° E) to Greece (38° N 21° E) (Fig. 1). The mesocosms were
95 set up in the lakes listed in Table 1. The mesocosms were closed systems (i.e. no direct
96 connection with the lake water column or bottom sediments) but were exposed to the same
97 climatic forcing as the lakes. The periphyton experiment presented here is part of a
98 comprehensive study of the effects of climate change on shallow lake ecosystems, which
99 started in May 2011 and continued until the end of October of the same year. Set-up and
100 sampling were standardized by a common protocol to ensure comparability between the
101 countries (Landkildehus et al. 2014). The present study lasted four weeks between 15 July and
102 15 August 2011, thus reflecting mid-summer growth conditions. The mesocosms included in
103 this experiment (8 in each country) consisted of 2.2 m deep cylindrical enclosures made of
104 fiberglass with a diameter of 1.2 m. The experimental treatment design comprised two
105 nutrient levels, resembling mesotrophic and eutrophic conditions. Manipulation of nutrient
106 levels was carried out using inorganic phosphate (P) [Na_2HPO_4] and nitrogen (N) [$\text{Ca}(\text{NO}_3)_2$]
107 at an N:P mass ratio of 1:20. Nutrients were added to the mesocosms at the beginning of the
108 experiment to get starting P and N concentrations of 25 $\mu\text{g L}^{-1}$ and 0.5 mg L^{-1} in the
109 mesotrophic (low loading) treatment and 200 $\mu\text{g L}^{-1}$ and 2 mg L^{-1} in the eutrophic (high
110 loading) treatment, respectively. Later, during the course of the experiment, monthly nutrient
111 additions amounted to 10.8 mg of P and 216 mg of N per mesocosm at low loading and 172
112 mg of P and 3440 mg of N at high loading (Landkildehus et al., 2014). These nutrient
113 additions took place after monthly sampling of the mesocosms. For each nutrient treatment,
114 four replicates were implemented in each country. During the mesocosm set-up in May 2011,
115 a 10 cm layer of sediments was added to all mesocosms (90% washed sand and 10% natural
116 sediment from oligotrophic local lakes). Subsequently, the mesocosms were filled with sieved

117 lake water (mesh size 500 μm) in all countries, except for Germany and the Czech Republic
118 where tap water was used because the lake water TP concentration was higher than the target
119 concentration of the low nutrient treatment (i.e. $> 25 \mu\text{g L}^{-1}$). The initial water level in each
120 mesocosms was 2 m. To ensure that naturally occurring phytoplankton, zooplankton, and
121 macroinvertebrate communities would emerge, the mesocosms were inoculated with plankton
122 and sediment samples, which were collected from five different local lakes covering a range
123 from oligotrophic to eutrophic conditions (Landkildehus et al. 2014). The mesocosm set-up
124 also included the addition of apical shoots of macrophytes (*Myriophyllum spicatum*). Six
125 adult planktivorous fish (length 2-4 cm, 3 males and 3 females to allow breeding) were
126 stocked in each enclosure at the beginning of the experiment. Three-spined sticklebacks
127 (*Gasterosteus aculeatus*) were used in all countries except of Greece where mosquito fish
128 (*Gambusia affinis*) were used. Both fish species are known to have similar diets (Offill and
129 Walton 1999; Simpson 2008). Dead fish were replaced during the experiment. The water of
130 the mesocosms was continuously circulated by using water pumps. A more detailed
131 description of the entire experimental set-up can be found in Landkildehus et al. (2014).

132

133 Variables measured

134 Periphyton growth over the experimental period was quantified based on the biomass
135 accumulation on artificial transparent polypropylene strips (2 strips, 16 x 2 cm) with a slightly
136 textured surface (IBICO®, Germany; Roberts et al. 2003). The strips were exposed at a water
137 depth of 0.5 m and kept 0.3 m away from the mesocosm walls facing south to prevent shading
138 from the walls, and the backsides of the strips were covered with adhesive tape.

139 After five weeks of colonization, the periphyton strips were gently lifted to the surface
140 to minimize disturbance and loss of periphyton mats. After removal of the adhesive tape from
141 the backside of the strips, these were immediately placed in round plastic tubes and

142 transported to the laboratory in a portable cooler box containing tap water to prevent the
143 samples from drying out.

144 For periphyton dry weight and chlorophyll *a* analysis, periphyton was scrubbed from
145 the strips using a soft toothbrush and suspended in a defined amount of filtered mesocosm
146 water (two cellulose acetate filters, diameter 50 mm, pore sizes 0.24 and 0.8 µm). Before
147 scrubbing, invertebrate grazers (mostly cladocerans and chironomids) were removed from the
148 strips using carbonated water (3-5 min exposure). After homogenization, aliquot subsamples
149 of each suspension were filtered onto two pre-weighed and pre-washed glassfiber filters
150 (Whatman GF/C, diameter 25 mm, pore size 0.7 µm) and dried at 105 °C for 12 h to analyze
151 periphyton dry weight. Ash-free dry weight was determined after combustion at 500 °C for 5
152 h. For chlorophyll *a* analyses, aliquot samples were filtered through glassfiber filters
153 (Whatman GF/F; 25 mm). Concomitantly with the periphyton harvest, water samples were
154 taken to determine concentrations of total phosphorus and total nitrogen, and phytoplankton
155 chlorophyll *a*. In each country, chlorophyll *a* (periphyton and phytoplankton), total
156 phosphorus, and total nitrogen concentrations were determined using the procedures
157 described in Landkildehus et al. (2014). Macrophyte plant volume inhabited (PVI %) was
158 calculated using the formula: $PVI (\%) = \% \text{ coverage} \times \text{average height} / \text{water depth}$, and
159 percent coverage and average height were visually estimated.

160 Mean air temperature for the experimental period was calculated from daily mean air
161 temperature data (hourly values). Air temperature and global radiation data were provided by
162 the Centre for Limnology of the Estonian University of Life Sciences, Leibniz-Institute of
163 Freshwater Ecology and Inland Fisheries (IGB), Czech Hydrometeorological Institute,
164 Turkish State Meteorology Service, and Hellenic National Meteorological Service.

165 Daily mean water temperature (24 hour averages of samples taken every 2 hours) was
166 measured on two occasions in July and August (11 July 2011 and 8 August 2011). The July–
167 August average water temperature values were used in the analysis (Table 2) as they

168 represented well the average temperate conditions for the experimental period, established by
 169 the mean air temperature. The close link between air temperature and surface water
 170 temperature in shallow lakes is well established in the literature (McCombie 1959;
 171 Livingstone and Lotter 1998; Mooij et al. 2008). At midday of the 24 h measurement events,
 172 profiles of photosynthetically active radiation (PAR) were taken at 0.1 m intervals from top to
 173 bottom. For each light profile and each concurrent light intensity measurement, an attenuation
 174 coefficient K_{di} (m^{-1}) was estimated based on the Beer-Lambert law:

$$K_{di} = \frac{\ln\left(\frac{I_i}{I_{i+1}}\right)}{z_{i+1} - z_i} \quad (1)$$

175 where I_i and I_{i+1} are PAR values at depth z_i and z_{i+1} , respectively. Values with $I_{i+1} > I_i$ were
 176 removed. K_d (m^{-1}) was then taken as the mean over all K_{di} . The attenuation coefficients from
 177 July and August were subsequently averaged. Mean and maximum available PAR at 0.5 m
 178 were calculated as: $I_{0.5(\text{mean or max})} = I_{0(\text{mean or max})} \exp(-0.5 K_d)$, where I_0 was set to average light
 179 hour PAR ($I_{0.5\text{mean}}$) or maximum PAR ($I_{0.5\text{max}}$) at the surface. Averages for the experimental
 180 period were calculated from daily $I_{0.5(\text{mean or max})}$ values. PAR was estimated from global
 181 radiation as $\text{PAR} = E \times \gamma \times 0.45$, where E is global radiation and $\gamma = 4.6$ is the mean photon
 182 flux in the wavelengths from 400 – 700 nm (Kirk 2010).
 183

184 At the end of the mesocosm experiment in November 2011, macroinvertebrates were
 185 sampled with the help of Kajak cores (diameter =52 mm) or an Ekman grab sampler.
 186 Subsequent identification and enumeration of snails were carried out to genus or species level.
 187 At the same time, all fish were captured and weighed. In Germany, three mesocosms (two in
 188 the high nutrient and one in the low nutrient treatment) sank during heavy storm events and
 189 were consequently omitted. A detailed description of the sampling procedure and processing
 190 can be found in Landkildehus et al. (2014).
 191

192 Data analyses

193 We analyzed the data using analysis of covariance (ANCOVA) to test for significant
194 differences in periphyton dry weight and chlorophyll *a* between the two nutrient treatments.
195 We did not test for the effect of nutrient treatment on periphyton ash free dry weight because
196 this variable was closely correlated with periphyton dry weight ($r^2= 0.91, p= 0001$).
197 ANCOVA was chosen because of the use of two discrete nutrient treatments (low and high)
198 in the experimental mesocosm set-up, these being considered factors in the analysis.
199 However, since the measured nutrient concentrations in the mesocosms showed considerable
200 variation within and among treatments (i.e. low and high), we confirmed the suitability of
201 applying ANCOVA by testing for significant differences in TP and TN concentrations
202 between treatments using one-way ANOVA.

203 The appropriateness of including potential covariates in the model was tested prior to
204 conducting the ANCOVA analysis. The candidate covariates were submerged macrophyte
205 PVI, snail abundance, fish biomass, phytoplankton chlorophyll *a* concentrations, mean PAR
206 at 0.5 m, and maximum PAR at 0.5 m. We used one-way ANOVA to test for significant
207 differences for each variable between the two nutrient treatments. The appropriateness of
208 including covariates was rejected if the factor nutrient treatment significantly affected a
209 particular variable. Furthermore, pairwise Pearson product moment correlation coefficients
210 were calculated between candidate covariates to ensure that selected covariates were not
211 strongly correlated (not reported). Based on the analysis above, water temperature, snail
212 abundances, and fish biomass were selected as covariates in the ANCOVA models. Because
213 submerged macrophyte PVI, phytoplankton chlorophyll *a* concentrations, mean PAR at 0.5
214 m, and maximum PAR at 0.5 m showed significant differences between nutrient treatments,
215 these variables were not included as covariates (see Fig. 2a-f). Snail abundance and fish
216 biomass were presumed to reflect the grazing pressure on periphyton. We assumed higher
217 periphyton grazing with higher snail abundance and lower periphyton grazing with increasing

218 fish biomass due to their higher predation on invertebrates (cascading effect) (Liboriussen et
219 al. 2005). A weak negative effect of fish on snails was found by regression analysis between
220 fish biomass and snail abundance ($b = 0.08$, $t(35) = 1.96$, $p = 0.06$) and logistic regression
221 analysis between fish biomass and presence and absence of snails ($b = -0.53$, $z(35) = -2.08$, p
222 $= 0.04$). Therefore, two alternative ANCOVA models (using either snail abundance or fish
223 biomass as a covariate) were analyzed for both periphyton dry weight and periphyton
224 chlorophyll *a*. All models contained nutrient treatment (i.e. high and low) as the main factor
225 and water temperature as a covariate. Although snail abundance data were used in this
226 analysis, it should be noted that snail species composition varied between countries. The
227 ANCOVA was executed using a Type III sums of squares method to account for the
228 unbalanced design in our experiment arising from the loss of three mesocosms in Germany.
229 Where necessary, data were either log or square root transformed to improve normality of the
230 residuals and meet the assumption of homogeneity of variance.

231 Additionally, to the ANCOVA analysis, differences in periphyton dry weight and
232 periphyton chlorophyll *a* between nutrient treatments were tested within each country. The
233 non-parametric Mann-Whitney U-test was considered appropriate to test for statistical
234 differences between treatments owing to the small sample size (i.e. $N=8$) in each country.
235 This test was not conducted for Germany due to the small sample size after losing three
236 mesocosms during the experiment (see above).

237 To aid the interpretation of the ANCOVA and to potentially further isolate the effect
238 of fish on periphyton biomass, an additional regression analysis was carried out between
239 water temperature-adjusted periphyton dry weight and chlorophyll *a* and fish biomass.
240 Adjustment involved calculating residuals of the regression equation of periphyton dry weight
241 vs. water temperature and periphyton chlorophyll *a* vs. water temperature, respectively. All
242 analyses were undertaken using STATISTICA 12 (StatSoft, Inc. USA) with a significance
243 threshold for all tests of $p \leq 0.05$.

244 **Results**

245 A clear temperature gradient was obtained by deploying mesocosms in five countries across
246 Europe simultaneously (Fig. 3a). Average air temperatures over the study period ranged from
247 17.0 °C in the Czech Republic to 27.3 °C in Greece. Established water temperatures were
248 strongly correlated with average air temperature ($R=0.97$, $p<0.001$) and were either equal or
249 slightly warmer than average air temperatures, which is consistent with a high heat storage
250 capacity of large water bodies (Table 2, Fig. S1). The measured average TP concentrations in
251 the low nutrient treatment were $20.1 \pm 6.9 \mu\text{g L}^{-1}$, while the measured TP concentrations in
252 the high nutrient treatment were $65.4 \pm 27.8 \mu\text{g L}^{-1}$ (Table 2). The mean TP concentration
253 difference between the two treatments was significant (ANOVA, $F_{1,36}= 81.57$, $p< 0.001$; Fig.
254 4a). Total nitrogen concentrations did not differ between the high and low nutrient treatments
255 ($F_{1,36}= 3.813$, $p= 0.059$, average of $1.46 \pm 1.05 \text{ mg L}^{-1}$ and $0.82 \pm 0.36 \text{ mg L}^{-1}$ for the high and
256 low nutrient treatment, respectively; Fig. 4b).

257 The summary statistics of all potential candidate covariates for the ANCOVA model
258 are presented in Table 2. Nutrient treatment had no significant effect on the candidate
259 covariates snail abundance and fish biomass (ANOVA, $F_{1,36}=0.042$, $p=0.839$; $F_{1,36}=0.11$,
260 $p=0.742$; $F_{1,36}=1.005$, $p=0.323$, respectively), but significant effects on macrophytes, water
261 column chlorophyll *a*, and mean and the maximum PAR were observed (Fig. 2a-g). Snails
262 were present in mesocosms in Estonia (*Valvata piscinalis*), the Czech Republic (*Lymnaea*
263 *stagnalis*), and Turkey (members of Planorbidae, Physidae, Lymnaeidae) but absent in
264 Germany and Greece.

265 The results of the ANCOVA analysis are summarized in Table 3. Overall, nutrient
266 treatment had no significant effect on either periphyton dry weight or chlorophyll *a*. Water
267 temperature was a significant covariate in all models, except for periphyton chlorophyll *a*
268 when snail abundance was included as a second covariate. Snail abundance was a significant
269 covariate in both models, for periphyton dry weight and chlorophyll *a*, respectively. Fish

270 biomass was a significant covariate for periphyton chlorophyll *a* but not for periphyton dry
271 weight.

272 The results of the Mann-Whitney U test showed that nutrient treatment did not have a
273 significant effect on periphyton dry weight (Fig. 3b) or chlorophyll *a*, except for periphyton
274 chlorophyll *a* in Greece (Fig. 3c). Periphyton dry weight was significantly correlated with
275 water temperature ($r^2=0.41$, $p=0.001$; Fig. 5a) and periphyton chlorophyll *a* ($r^2=0.28$,
276 $p=0.001$; Fig. 5c). Fish biomass showed a weak relationship with temperature-adjusted
277 periphyton dry weight ($r^2=0.1$, $p=0.10$) (Fig. 5b) and a strongly significant relationship with
278 temperature-adjusted periphyton chlorophyll *a* ($r^2=0.45$, $p<0.001$) (Fig. 5d).

279

280 **Discussion**

281 The present pan-European lake mesocosm experiment provided evidence that increasing
282 water temperature can lead to increased development of summer periphyton biomass. Nutrient
283 enrichment had no significant effect on periphyton biomass, probably due to very low nutrient
284 limitation levels for periphyton. Indirect top-down effects of fish emerged as an important
285 factor controlling periphyton biomass and appeared to be independent of water temperature.
286 In addition, snails, when present, appeared to have a negative effect on periphyton chlorophyll
287 *a*.

288 Our results showed that periphyton biomass (measured as dry weight) was
289 significantly and positively correlated with water temperature in the 20-28°C temperature
290 range. Given the projected rise in global air and water temperatures (IPCC 2013), our results,
291 therefore, suggest that summer periphyton biomass is likely to increase in the future. Our
292 results contradict those of Hansson (1992) who found that temperature was of minor
293 importance for periphyton biomass. However, his study was conducted along a much larger
294 productivity gradient (Swedish and Antarctic lakes) ranging from extremely low (meltwater
295 lakes) to highly productive lakes. In our study, total phosphorus concentrations covered a

296 smaller range (20-65 $\mu\text{g L}^{-1}$) and had no significant effect on periphyton biomass. Others have
297 found a unimodal relationship between periphyton biomass and total phosphorus, peaking at
298 39 $\mu\text{g L}^{-1}$ (Lalonde and Downing 1991) or between 60-200 $\mu\text{g L}^{-1}$ (Liboriussen and Jeppesen
299 2006). The concentrations of dissolved reactive silicon were mostly above limiting levels (0.5
300 mg L^{-1}) in the low and high nutrient treatments of the Czech Republic ($0.9 \pm 0.1 \text{ mg L}^{-1}$ and
301 $0.3 \pm 0.1 \text{ mg L}^{-1}$) and Germany ($1.8 \pm 1.5 \text{ mg L}^{-1}$ and $1.0 \pm 1.2 \text{ mg L}^{-1}$). However, data are
302 lacking for the other countries. Light levels ranged on average between 39 and 408 μmol
303 $\text{photons m}^{-2} \text{ s}^{-1}$ and were always above the minimum light requirement for growth of
304 microalgae (1-10 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) given by Sand-Jensen & Borum (1991). Therefore,
305 light limitation was unlikely in our study.

306 Furthermore, our results are different from those obtained in a pan-European study by
307 Bécarea et al. (2008), covering a temperature range of 17.7-29°C. They found higher
308 periphyton chlorophyll *a* in northern lakes than in southern lakes and explained this by a
309 stronger shading effect by phytoplankton on periphyton in southern lakes. The phytoplankton
310 chlorophyll *a* concentrations in their study were generally higher (40-564 $\mu\text{g L}^{-1}$) than in our
311 study (2-53 $\mu\text{g L}^{-1}$), even if age of periphyton is comparable. The same applies for the
312 maximum periphyton chlorophyll *a* concentration (84 mg m^{-2}), which with was five times
313 higher than in our study (16 mg m^{-2}). Furthermore, they found nutrient concentrations to be an
314 important driver (tested at six levels of $\text{NO}_3^{-}\text{-N}$ and $\text{PO}_4^{3-}\text{-P}$ up to 100 mg L^{-1} and 10 mg L^{-1} ,
315 respectively). In contrast, a positive top-down effect of fish on periphyton biomass was found
316 in our study, which was indicated by the significant positive relation recorded between fish
317 biomass and periphyton chlorophyll *a* in both ANCOVA and regression analysis. This result
318 was probably due to the prevailing top-down control by fish of periphyton-scraping non-snail
319 invertebrates, as suggested by Jones and Sayer (2003) and Danger et al. (2008). Körner and
320 Dugdale (2003) showed a switch of fish to periphyton-scraping invertebrates at low
321 zooplankton abundance. In our experiment, in all countries except Greece, we used

322 sticklebacks, a bottom-feeder that essentially feeds on plankton and benthic prey (Sánchez-
323 González et al. 2001). However, periphyton biomass was highest in Greece where
324 mosquitofish were used instead of sticklebacks. Although both species feed mainly on the
325 same food items (planktonic and littoral zooplankton, chironomid larvae) (Offill and Walton
326 1999; Simpson 2008), we cannot rule out the potential occurrence of confounding factors in
327 the cascading effects of different fish species in the mesocosms.

328 Snail abundance had a significant effect on periphyton biomass in our study, but snails
329 were absent in Germany and Greece. Snails may have directly scraped periphyton as known
330 from various other studies (e.g. Brönmark 1989) and thus contributed to the low periphyton
331 biomass observed in the Czech Republic, Estonia, and Turkey. Nutrient recycling from snail
332 faeces and excreta might also have increased nutrient availability for periphyton in the low
333 nutrient treatments and contributed to the lack of differences in periphyton biomass compared
334 to the high nutrient treatments in these countries (Liess and Haglund 2007). Periphyton
335 biomass was, however, also low in Germany without the presence of snails. Given the size of
336 sticklebacks, top-down effects of fish on snails (Brönmark et al. 1992) seem unlikely (snail
337 size: 5 mm – 7 cm). Yet, snail abundance tended to be lower if fish were present.

338 In general, in Mediterranean shallow lakes fish seem to exert strong trophic cascading
339 effects due to dominance by frequently spawning omnivores and benthivores and absence of
340 efficient piscivores (Beklioglu et al. 2007; Papastergiadou et al. 2010). Gyllström et al. (2005)
341 found that the ratio between prey and predators and fish:zooplankton biomass increased from
342 northern to southern Europe, while the zooplankton:phytoplankton biomass ratio decreased.
343 The absence of large-bodied zooplankton due to strong fish predation seems to be the reason
344 for lack of phytoplankton control, and a similar mechanism may explain the lack of top-down
345 control of periphyton by scraping invertebrates in Greece. In contrast, fish biomass in the
346 Turkish mesocosms was low, which might explain the low periphyton biomass despite
347 warmer conditions. Therefore, the overall effect of temperature on periphyton seems to

348 depend on nutrient level (Hansson 1992; Liboriussen et al. 2005; Trochine et al. 2014), fish
349 abundance and composition (some being periphyton grazers; Gonzáles-Bergonzoni et al.
350 2012), and the strength of the cascading effects of fish on invertebrate periphyton grazers
351 (Cao et al. 2014; Meerhoff et al. 2007).

352 In conclusion, our results indicate a stimulating effect of water temperature on summer
353 periphyton biomass. Due to non-limiting nutrient levels and low differences between the
354 treatments, no significant effect of nutrient loading on periphyton biomass was observed.
355 However, apart from temperature, direct and indirect top-down control of snails and fish
356 proved to be important factors for explaining a significant amount of variation in periphyton
357 biomass.

358

359 **Acknowledgements**

360 This project was primarily supported by the EU FP-7 Theme 6 project REFRESH
361 (Adaptive strategies to Mitigate the Impacts of Climate Change on European Freshwater
362 Ecosystems, Contract No.: 244121). EJ, MS, and FL were also supported by ‘CLEAR’ (a
363 Villum Kann Rasmussen Centre of Excellence project), CRES, and CIRCE. In addition, JH
364 and MS obtained support from the MSMT CR (project No. 425 7E11059)) and NF was
365 supported by TÜBİTAK 2211 Scholarship programme, and the study was partly supported by
366 TÜBİTAK (ÇAYDAĞ110Y125). AM was supported by a PhD426 stipend from the Egyptian
367 Government. Support also came from the Estonian Ministry of Education (SF 0170011s08)
368 and Estonian Science Foundation grants 8729 and 9102, and EP, KS, and CP were supported
369 by grants from the University of Patras. We want to thank Arvo Tuvikene, Tõnu Feldmann,
370 Helen Agasild, Anu Kisand, Katrit Karus, Şeyda Erdoğan, Tuba Bucak, Eti Levi, Gizem
371 Bezirci, Arda Özen, Engin Bilgen, Tasos Samiotis, and Yannis Nikolopoulos for help with
372 setting up the experiment, nursing the imported fish, and undertaking the regular sampling.
373 The local fishermen in Lake Lysimachia are acknowledged for keeping the mesocosm

374 construction safe. The Erken Field laboratory and the technicians of the Department of
375 Aquatic Sciences and Assessment (SLU) also provided invaluable help. The IGB provided
376 additional funding for the experimental set-up. We thank the staff at IGB for their tremendous
377 technical support during the set-up of the experiments, sampling, and sample processing;
378 especially Thomas Hintze. We also thank Armaplast Co. for constructing the mesocosms and
379 ensuring their safe transport to all the experimental sites. Anne Mette Poulsen's linguistic
380 improvements are acknowledged. The map of Europe in Figure 1 was provided by Natural
381 Earth and customized with QGIS 2.0.1-Dufour.

382 **References**

- 383 Adrian R, O'Reilly CM, Zagarese H, Baines SB, Hessen DO, Keller W, Livingstone DM,
384 Sommaruga R, Straile D, Van Donk E, Weyhenmeyer GA, Winder M (2009) Lakes as
385 sentinels of climate change. *Limnol Oceanogr* 54:2283-2297.
386 doi:10.4319/lo.2009.54.6_part_2.2283
- 387 Adrian R, Wilhelm S, Gerten D (2006) Life-history traits of lake plankton species may govern
388 their phenological response to climate warming. *Global Change Biol* 12:652-661.
389 doi:10.1111/j.1365-2486.2006.01125.x
- 390 Bécares E, Gomá J, Fernández-Aláez M, Fernández-Aláez C, Romo S, Miracle M, Ståhl-
391 Delbanco A, Hansson L-A, Gyllström M, Bund W, Donk E, Kairesalo T, Hietala J,
392 Stephen D, Balayla D, Moss B (2008) Effects of nutrients and fish on periphyton and
393 plant biomass across a European latitudinal gradient. *Aquat Ecol* 42:561-574.
394 doi:10.1007/s10452-007-9126-y
- 395 Beklioğlu M, Romo S, Kagalou I, Quintana X, Bécares E (2007) State of the art in the
396 functioning of shallow Mediterranean lakes: workshop conclusions. *Hydrobiologia*
397 584:317-326. doi:10.1007/s10750-007-0577-x
- 398 Brönmark C (1989) Interactions between epiphytes, macrophytes and freshwater snails: a
399 review. *J Moll Stud* 55:299-311. doi: 10.1093/mollus/55.2.299
- 400 Brönmark C, Klosiewski SP, Stein RA (1992) Indirect effects of predation in a freshwater
401 benthic food chain. *Ecology* 73:1662-1674. doi: 10.2307/1940018
- 402 Cao Y, Li W, Jeppesen E (2014). The response of two submerged macrophyte and periphyton
403 to elevated temperature at high nutrient level: a microcosm approach. *Hydrobiologia*
404 738:49-59. doi: 10.1007/s10750-014-1914-5
- 405 Danger M, Lacroix G, Oumarou C, Benest D, Meriguet J (2008) Effects of food-web structure
406 on periphyton stoichiometry in eutrophic lakes: a mesocosm study. *Freshw Biol*
407 53:2089-2100. doi:10.1111/j.1365-2427.2008.02031.x

408 González-Bergonzoni I, Meerhoff M, Davidson T, Teixeira-de Mello F, Baattrup-Pedersen A,
409 Jeppesen E (2012) Meta-analysis shows a consistent and strong latitudinal pattern in
410 fish omnivory across ecosystems. *Ecosystems* 15:492-503. doi:10.1007/s10021-012-
411 9524-4

412 Gyllström M, Hansson LA, Jeppesen E, García-Criado F, Gross E, Irvine K, Kairesalo T,
413 Kornijow R, Miracle MR, Nykänen M, Nöges T, Romo S, Stephen D, Van Donk E,
414 Moss B (2005) The role of climate in shaping zooplankton communities of shallow
415 lakes. *Limnol Oceanogr* 50:2008-2021. doi:10.4319/lo.2005.50.6.2008

416 Hansson LA (1992) Factors regulating periphytic algal biomass. *Limnol Oceanogr* 37:322-
417 328.

418 IPCC (2013) Summary for policymakers. In: Stocker TF, Qin D, Plattner GK, Tignor M,
419 Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (eds) *Climate change*
420 *2013: the physical science basis. Contribution of working group I to the fifth*
421 *assessment report of the intergovernmental panel on climate change. Cambridge*
422 *University Press, Cambridge, United Kingdom and New York, NY, USA*

423 Jeppesen E, Jensen J, Søndergaard M, Lauridsen T, Pedersen L, Jensen L (1997) Top-down
424 control in freshwater lakes: the role of nutrient state, submerged macrophytes and
425 water depth. *Hydrobiologia* 342/343:151-164. doi:10.1023/a:1017046130329

426 Jones JI, Sayer CD (2003) Does the fish-invertebrate-periphyton cascade precipitate plant loss
427 in shallow lakes? *Ecology* 84:2155-2167. doi:10.1890/02-0422

428 Kirk JTO (2010) *Light and photosynthesis in aquatic ecosystems. 3rd Edition. Cambridge*
429 *University Press*

430 Körner S, Dugdale T (2003) Is roach herbivory preventing re-colonization of submerged
431 macrophytes in a shallow lake? *Hydrobiologia* 506/509:497-501. doi:
432 10.1023/B:HYDR.0000008561.67513.ec

433 Lalonde S, Downing JA (1991) Epiphyton biomass is related to lake trophic status, depth, and
434 macrophyte architecture. *Can J Fish Aquat Sci* 48:2285-2291. doi:10.1139/f91-268

435 Landkildehus F, Søndergaard M, Beklioglu M, Adrian R, Angeler DG, Hejzlar J,
436 Papastergiadou E, Zingel P, Çakiroğlu Aİ, Scharfenberger U, Drakare S, Nöges T,
437 Šorf M, Stefanidis K, Tavşanoğlu ÜN, Trigal C, Mahdy A, Papadaki C, Tuvikene L,
438 Kernan M, Jeppesen E (2014) Climate change effects on shallow lakes: Design and
439 preliminary results of a cross-European climate gradient mesocosm experiment.
440 *Estonian J Ecol* 63:71-89. doi: 10.3176/eco.2014.2.02

441 Liboriussen L, Jeppesen E, Bramm M, Lassen M (2005) Periphyton-macroinvertebrate
442 interactions in light and fish manipulated enclosures in a clear and a turbid shallow
443 lake. *Aquat Ecol* 39:23-39. doi:10.1007/s10452-004-3039-9

444 Liboriussen L, Jeppesen E (2006) Structure, biomass, production and depth distribution of
445 periphyton on artificial substratum in shallow lakes with contrasting nutrient
446 concentrations. *Freshw Biol* 51:95-109. doi: 10.1111/j.1365-2427.2005.01481.x

447 Liess A, Haglund AL (2007) Periphyton responds differentially to nutrients recycled in
448 dissolved or faecal pellet form by the snail grazer *Theodoxus fluviatilis*. *Freshw Biol*
449 52:1997-2008. doi:10.1111/j.1365-2427.2007.01825.x

450 Livingstone DM, Lotter AF (1998) The relationship between air and water temperatures in
451 lakes of the Swiss Plateau: a case study with palaeolimnological implications. *J*
452 *Paleolimnol* 19:181-198. doi:10.1023/A:1007904817619

453 McCombie AM (1959) Some relations between air temperatures and the surface water
454 temperature of lakes. *Limnol Oceanogr* 4:252-258. doi:10.4319/lo.1959.4.3.0252

455 McKee D, Hatton K, Eaton JW, Atkinson D, Atherton A, Harvey I, Moss B (2002) Effects of
456 simulated climate warming on macrophytes in freshwater microcosm communities.
457 *Aquat Bot* 74:71-83. doi:10.1016/S0304-3770(02)00048-7

458 Meerhoff M, Clemente JM, De Mello FT, Iglesias C, Pedersen AR, Jeppesen E (2007) Can
459 warm climate-related structure of littoral predator assemblies weaken the clear water
460 state in shallow lakes? *Glob Change Biol* 13:1888-1897. doi:10.1111/j.1365-
461 2486.2007.01408.x

462 Meerhoff M, Teixeira-de Mello F, Kruk C, Alonso C, González-Bergonzoni I, Pacheco JP,
463 Lacerot G, Arim M, Beklioglu M, Brucet S, Goyenola G, Iglesias C, Mazzeo N,
464 Kosten S, Jeppesen E (2012) Environmental warming in shallow lakes: a review of
465 potential changes in community structure as evidenced from space-for-time
466 substitution approaches. *Advances in Ecological Research*, vol 46. Academic Press, pp
467 259-349. doi:10.1016/B978-0-12-396992-7.00004-6

468 Mooij WM, Hülsmann S, De Senerpont Domis L, Nolet B, Bodelier PE, Boers PM, Pires LM,
469 Gons H, Ibelings B, Noordhuis R, Portielje R, Wolfstein K, Lammens ERR (2005)
470 The impact of climate change on lakes in the Netherlands: a review. *Aquat Ecol*
471 39:381-400. doi:10.1007/s10452-005-9008-0

472 Mooij WM, De Senerpont Domis L, Hülsmann S (2008) The impact of climate warming on
473 water temperature, timing of hatching and young-of-the-year growth of fish in shallow
474 lakes in the Netherlands: *J Sea Res* 60:32-43. doi:10.1016/j.seares.2008.03.002

475 Offill YA, Walton WE (1999) Comparative efficacy of the threespine stickleback
476 (*Gasterosteus aculeatus*) and the mosquitofish (*Gambusia affinis*) for mosquito
477 control. *J Am Mosq Control Assoc* 15:380-390.

478 Papastergiadou E, Kagalou I, Stefanidis K, Retalis A, Leonardos I (2010) Effects of
479 anthropogenic influences on the trophic state, land uses and aquatic vegetation in a
480 shallow Mediterranean lake: implications for restoration. *Water Resour Manage*
481 24:415-435. doi:10.1007/s11269-009-9453-y

482 Patrick DA, Boudreau N, Bozic Z, Carpenter GS, Langdon DM, LeMay SR, Martin SM,
483 Mourse RM, Prince SL, Quinn KM (2012) Effects of climate change on late-season

484 growth and survival of native and non-native species of watermilfoil (*Myriophyllum*
485 spp.): Implications for invasive potential and ecosystem change. *Aquat Bot* 103:83-88.
486 doi:10.1016/j.aquabot.2012.06.008

487 Phillips GL, Eminson D, Moss B (1978) A mechanism to account for macrophyte decline in
488 progressively eutrophicated freshwaters. *Aquat Bot* 4:103-126. doi:10.1016/0304-
489 3770(78)90012-8

490 Roberts E, Kroker J, Körner S, Nicklisch A (2003) The role of periphyton during the re-
491 colonization of a shallow lake with submerged macrophytes. *Hydrobiologia* 506-
492 509:525-530. doi:10.1023/B:HYDR.0000008560.73832.1c

493 Sánchez-González S, Ruiz-Campos G, Contreras-Balderas S (2001) Feeding ecology and
494 habitat of the threespine stickleback, *Gasterosteus aculeatus microcephalus*, in a
495 remnant population of northwestern Baja California, México. *Ecol Freshw Fish*
496 10:191-197. doi:10.1034/j.1600-0633.2001.100401.x

497 Sand-Jensen K, Borum J (1991) Interactions among phytoplankton, periphyton, and
498 macrophytes in temperate freshwaters and estuaries. *Aquat Bot* 41:137-175.
499 doi:10.1016/0304-3770(91)90042-4

500 Scheffer M, Hosper SH, Meijer ML, Moss B, Jeppesen E (1993) Alternative equilibria in
501 shallow lakes. *Trends Ecol Evol* 8:275-279. doi: 10.1016/0169-5347(93)90254-M.

502 Seebens H, Einsle U, Straile D (2009) Copepod life cycle adaptations and success in response
503 to phytoplankton spring bloom phenology. *Glob Change Biol* 15:1394-1404.
504 doi:10.1111/j.1365-2486.2008.01806.x

505 Shurin JB, Clasen JL, Greig HS, Kratina P, Thompson PL (2012) Warming shifts top-down
506 and bottom-up control of pond food web structure and function. *Phil T R Soc B-Biol*
507 *Sci* 367 (1605):3008-3017. doi:10.1098/rstb.2012.0243

508 Simpson, N.P. 2008. The potential impact on mosquito larvae by threespine stickleback
509 (*Gasterosteus aculeatus*) and mosquitofish (*Gambusia affinis*) in four constructed

510 wetlands. Thesis (M.S.) – Humboldt State University, Fisheries: Wastewater
511 Utilization, 82 pp.

512 Tarkowska-Kukuryk M, Mieczan T (2012) Effect of substrate on periphyton communities and
513 relationships among food web components in shallow hypertrophic lake. *J Limnol*
514 71:279-290. doi: 10.4081/jlimnol.2012.e30

515 Teixeira-de Mello F, Meerhoff M, Pekcan-Hekim Z, Jeppesen E (2009) Substantial
516 differences in littoral fish community structure and dynamics in subtropical and
517 temperate shallow lakes. *Freshw Biol* 54:1202-1215. doi:10.1111/j.1365-
518 2427.2009.02167.x

519 Trochine C, Guerrieri M, Liboriussen L, Lauridsen TL, Jeppesen E (2014) Effects of nutrient
520 loading, temperature regime and grazing pressure on nutrient limitation of periphyton
521 in experimental ponds. *Freshw Biol* 59:905-917. doi: 10.1111/fwb.12314

522 Wagner C, Adrian R (2009) Exploring lake ecosystems: hierarchy responses to long-term
523 change? *Glob Change Biol* 15:1104-1115. doi:10.1111/j.1365-2486.2008.01833.x
524

525 **Table 1:** Lake mesocosm experiment – basic information about sites and mean conditions during the entire experimental period from May to
 526 October (modified from Landkildehus et al. 2014).

Experimental site	Coordinates	Climate	Altitude (m a.s.l)	No. of mesocosms	Total precipitation (mm)	Mean air temperature (°C)
Czech Republic, Vodňany	49°09'14"N; 14°10'11"E	Transient maritime/continental	395	8	401	15.3
Germany, Müggelsee	52°26'0" N; 13°39'0" E	Transient maritime/continental	32	5	431	16.9
Estonia, Võrtsjärv	58°12'17" N; 26°06'16" E	Boreal	35	8	298	14.4
Turkey, ODTU-DSI Golet	39°52'38" N; 32°46'32" E	Transient continental/Mediterranean	998	8	223	18.8
Greece, Lysimachia	38°33'40" N; 21°22'10" E	Mediterranean	16	8	252	23.9

527 **Table 2:** Mid-summer (i.e. 15 July – 15 August) means (\pm standard deviation) of selected variables in mesocosms with different nutrient
528 treatments (high and low) in five European countries. All means are treatment means apart from mean air temperature, which is averaged
529 over daily mean values.

Countries	Czech Republic		Germany		Estonia		Turkey		Greece	
	High	Low	High	Low	High	Low	High	Low	High	Low
Periphyton AFDW (g m ⁻²)	0.5±0.3	0.5±0.2	1.5±0.2	1.7±0.2	2.2±1.6	1.0±0.2	0.9±0.8	0.3±0.3	9.2±4.1	6.0±1.6
Periphyton chl <i>a</i> (mg m ⁻²)	5.0±3.4	2.8±1.0	0.9±0.1	0.7±0.0	0.1±0.0	0.04±0.0	0.4±0.4	0.1±0.1	16.8±5.7	3.1±0.9
TP (µg L ⁻¹)	84.6±31.3	15.2±3.2	40.0±5.7	25.0±8.5	45.0±9.6	14.0±0.8	65.7±35.9	19.9±4.6	79.0±17.2	29.0±2.9
TN (mg L ⁻¹)	0.8±0.3	0.9±0.31	3.4±0.5	0.6±0.1	1.4±0.2	0.8±0.0	0.7±0.1	0.4±0.1	1.1±0.4	1.4±0.2
Fish biomass (g m ⁻²)	1.3±0.2	0.9±0.2	1.4±0.5	0.9±0.2	0.0±0.0	0.0±0.0	0.3±0.4	0.2±0.2	0.7±0.1	0.7±0.7
Phytoplankton chl <i>a</i> (µg L ⁻¹)	5.4±2.1	5.7±1.3	72.5±11.0	6.9±1.6	20.4±12.4	13.4±3.6	9.8±3.4	2.2±1.1	52.5±61.7	7.0±8.1
Mean PAR at 0.5 m (µmol photons m ⁻² s ⁻¹)	146.3±17.7	178.8± 21.6	125.7±30.4	119.7±13.8	140.9±13.1	178.4±21.6	304.4±68.5	408.4 ± 55.5	38.6 ±11.7	218.6± 53.3
Max PAR at 0.5 m (µmol photons m ⁻² s ⁻¹)	447.3±54.1	546.6±66.0	265.0±64.0	252.3±29.1	313.2±29.1	396.4±48.0	614.0±138.2	823.9±111.9	70.9±21.4	401.9±98.0
Submerged macrophytes (% plant volume inhabited)	0.0±0.0	1.0±1.0	0.0±0.0	6.6±0.4	11.9±12.9	10.2±10.3	4.8±5.6	8.2±3.3	2.4±2.1	10.8±2.1
Snail abundance (individuals m ⁻²)	2.5±2.7	0.08±0.2	0.0±0.0	0.0±0.0	1.4±1.6	5.2±2.5	0.3±0.2	0.5±0.2	0.0±0.0	0.0±0.0
Mean water temperature (°C)	20.0 ±0.2	20.0 ±0.2	20.8 ±0.0	20.8 ±0.1	22.6 ±0.2	22.4 ±0.1	25.1 ±0.1	25.1 ±0.1	28.3 ±0.1	28.3 ±0.1
Mean air temperature (°C)	17.0 ± 2.4		18.0 ±1.9		19.2 ±3.2		25.8 ±2.5		27.3 ±1.4	

530 **Table 3:** Summary of results from the ANCOVA models testing for the effect of nutrient
 531 treatments on periphyton dry weight and periphyton chlorophyll *a*. Water temperature, snail
 532 abundance, and fish biomass were used as covariates in the models. Bold *p* values denote
 533 significant effects.

Dependent variable	Effect	SS	df	F	<i>p</i>
Periphyton dry weight	Intercept	6.561	1	13.386	<0.001
	Water temperature	9.415	1	19.211	<0.001
	Snail abundance	4.782	1	9.756	0.003
	Nutrient treatment	0.020	1	0.041	0.84
	Error	16.173	33		
Periphyton dry weight	Intercept	10.369	1	17.983	<0.001
	Water temperature	14.803	1	25.674	<0.001
	Fish biomass	1.927	1	3.342	0.076
	Nutrient treatment	0.003	1	0.005	0.942
	Error	19.028	33		
Periphyton chlorophyll <i>a</i>	Intercept	0.214	1	0.304	0.585
	Water temperature	0.045	1	0.065	0.801
	Snail abundance	7.089	1	10.096	0.003
	Nutrient treatment	1.343	1	1.912	0.176
	Error	23.173	33		
Periphyton chlorophyll <i>a</i>	Intercept	4.179	1	8.543	0.006
	Water temperature	2.94	1	6.011	0.019
	Fish biomass	14.121	1	28.869	<0.001
	Nutrient treatment	0.279	1	0.57	0.456
	Error	16.142	33		

534 **Figure captions**

535 **Fig. 1.** Map of Europe showing the five experimental locations: Estonia (Võrtsjärv), Germany
536 (Müggelsee), Czech Republic (Vodňany), Turkey (ODTÜ-DSİ Gölet), and Greece
537 (Lysimachia).

538

539 **Fig. 2.** Box and whisker plots representing the median values of (a) macrophyte plant volume
540 inhabited (PVI), (b) snail abundance, (c) fish biomass, (d) water column chlorophyll *a*
541 concentrations, (e) mean PAR measurements between July and August, and (f) maximum
542 PAR measurements between July and August for each nutrient treatment (high and low) in the
543 mesocosm experiments conducted in five European countries. Horizontal lines denote the
544 medians, boxes denote the 25th and 75th percentile, whiskers denote non-outlier range, circles
545 are outliers, and the asterisks are extreme values. *P* values were derived from a one-way
546 ANOVA to test for significant differences between nutrient treatments.

547

548 **Fig. 3.** Box and whisker plots representing the median values of (a) water temperature, (b)
549 periphyton dry weight, and (c) periphyton chlorophyll *a* content for each nutrient treatment
550 (high and low) in five European countries. Horizontal lines denote the medians, boxes denote
551 the 25th and 75th percentile, and whiskers denote non-outlier range. Asterisk indicates
552 significant differences between nutrient treatments based on Mann-Whitney U test at $p \leq 0.05$.

553

554 **Fig. 4.** Box and whisker plots representing the median values of (a) total phosphorus
555 concentrations and (b) total nitrogen concentrations for each nutrient treatment (high and low)
556 in the mesocosm experiments conducted in five European countries. Horizontal lines denote
557 the medians, boxes denote the 25th and 75th percentile, the whiskers denote non-outlier
558 range, circles are outliers. *P* values were derived from a one-way ANOVA to test for
559 significant differences between nutrient treatments.

560

561 **Fig. 5.** Relationship between (a) periphyton dry weight (DW) and water temperature (WT),
562 (b) periphyton dry weight (DW) adjusted for water temperature and fish biomass (g m^{-2}), (c)
563 periphyton chlorophyll *a* (chl *a*) and periphyton dry weight (DW), and (d) periphyton
564 chlorophyll *a* (chl *a*) adjusted for water temperature (WT) and fish biomass (g m^{-2}) in
565 mesocosm experiments in five European countries. Only significant *p*-values were included.

566



567

568

569

570

571

572

573

574

575

576


577

578

579

580


Fig. 1



581
582

583

584 Fig. 2



585

586

587

588

589

590


591

592

593

594

Fig. 3



595

596

597

598

599

600

601


602

603

604

605 Fig. 4

606



608

609

610

611

612

613

614

615 Fig. 5

616

617 **Supplementary material**

618

619 Figure S1. Comparison of mean daily air temperature values at study sites, with established
620 mean water temperatures (dashed horizontal lines) for the experimental period. Daily mean air
621 temperatures are based on hourly air temperature values. Average water temperature is based
622 on two daily mean values (24 hour averages of samples taken every two hours) measured on
623 11 July and 8 August 2011. CZ = Czech Republic, EE = Estonia, GE = Germany, GR =
624 Greece, and TR = Turkey.

625

626

